

# ANALYSIS & PDE

Volume 10      No. 8      2017

JONG-GUK BAK, JUNGIIN LEE AND SANGHYUK LEE

**BILINEAR RESTRICTION ESTIMATES  
FOR SURFACES OF CODIMENSION BIGGER THAN 1**





## BILINEAR RESTRICTION ESTIMATES FOR SURFACES OF CODIMENSION BIGGER THAN 1

JONG-GUK BAK, JUNGJIN LEE AND SANGHYUK LEE

In connection with the restriction problem in  $\mathbb{R}^n$  for hypersurfaces including the sphere and paraboloid, the bilinear (adjoint) restriction estimates have been extensively studied. However, not much is known about such estimates for surfaces with codimension (and dimension) larger than 1. In this paper we show sharp bilinear  $L^2 \times L^2 \rightarrow L^q$  restriction estimates for general surfaces of higher codimension. In some special cases, we can apply these results to obtain the corresponding linear estimates.

### 1. Introduction and statement of results

For a smooth hypersurface  $S$  such as the sphere or paraboloid in  $\mathbb{R}^n$ ,  $n \geq 3$ , the  $L^p$ - $L^q$  boundedness of the (adjoint) restriction operator (or the extension operator)  $\widehat{f}d\sigma$  has been extensively studied since the late 1960s. Here  $d\sigma$  denotes the induced Lebesgue measure on  $S$ . Specifically, when  $S$  is the sphere, it was conjectured by E. M. Stein [1993] that  $\widehat{f}d\sigma$  should map  $L^p(S)$  boundedly to  $L^q(\mathbb{R}^n)$ , precisely when  $q \geq p'(n+1)/(n-1)$  and  $q > 2n/(n-1)$ . Since then, a large amount of literature has been devoted to this problem. Over the last couple of decades, the bilinear and multilinear approaches have proven to be quite effective, and through them substantial progress has been made. We refer the reader to [Bennett et al. 2006; Bourgain and Guth 2011; Guth 2016] for the most recent developments.

On the other hand, when the dimension of the manifold is 1, namely, when the associated surface is a curve, the restriction estimate is by now fairly well understood [Bak et al. 2002; 2009; 2013; Stovall 2016].

However, not much is known about the intermediate cases, namely, when the codimension  $k$  of the manifold is between 1 and  $n-1$ . The restriction problem for quadratic surfaces of codimension  $k \geq 2$  was first studied by Christ [1982] and Mockenhaupt [1996]. They also considered the problem in a more general setting and found some necessary conditions on the curvature and codimension of the surface. For some surfaces they also established the optimal  $L^2 \rightarrow L^q$  linear estimates, which may be regarded as generalizations of the Stein–Tomas restriction theorem; see also [Banner 2002]. Although there are some known cases in which the  $L^p$ - $L^q$  boundedness is completely characterized, see for example [Bak and Ham 2014; Bak and Lee 2004; Oberlin 2005], for most surfaces with codimension bigger than 1, the current state of the restriction problem is hardly beyond that of the Stein–Tomas theorem.

---

J. Lee is supported in part by NRF grant No. 2017R1D1A1B03036053 (Republic of Korea). S. Lee is supported by NRF grant No. 2015R1A2A2A05000956 (Republic of Korea). J. Bak was supported in part by the Basic Science Research Institute (BSRI), POSTECH.

MSC2010: 42B15, 42B20.

Keywords: Fourier transform of measures, complex surfaces, Fourier restriction estimates.

In this paper, we are concerned with restriction estimates for surfaces of codimension  $k \geq 2$ . To be more specific, let us set  $k \geq 1$  and  $I = [-1, 1]$ . Let  $\Phi : I^d \rightarrow \mathbb{R}^k$  be a smooth function given by

$$\Phi(\xi) = (\varphi_1(\xi), \varphi_2(\xi), \dots, \varphi_k(\xi)).$$

The adjoint restriction operator (the extension operator)  $E = E_\Phi$  for the surface  $(\xi, \Phi(\xi)) \in \mathbb{R}^d \times \mathbb{R}^k$  is defined by

$$Ef(x, t) = \int_{I^d} e^{2\pi i(x \cdot \xi + t \cdot \Phi(\xi))} f(\xi) d\xi, \quad (x, t) \in \mathbb{R}^d \times \mathbb{R}^k.$$

Specific examples of such operators with  $2 \leq k \leq d - 2$  can be found in [Bak and Ham 2014; Bak and Lee 2004; Christ 1982; Mockenhaupt 1996; Oberlin 2005]. (Also, see Section 5.)

There are some classes of surfaces for which the optimal  $L^2$ - $L^q$  boundedness of  $E$  is well understood. In fact, using a Knapp-type example it is easy to see that  $E$  may be bounded from  $L^p$  to  $L^q$  only if  $(d + 2k)/q \leq d(1 - 1/p)$ . Hence, the best possible  $L^2$ - $L^q$  bound is that for  $q = 2(d + 2k)/d$ . Christ [1982] and Mockenhaupt [1996] showed that this is true for a class of surfaces satisfying a suitable curvature condition. In particular, let  $M$  be a linear map from  $\mathbb{R}^k$  to the space of  $d \times d$  symmetric matrices and suppose that  $\int_{S^{k-1}} |\det M(t)|^{-\gamma} d\sigma(t) < \infty$  for  $\gamma = k/d$ . Then it was proven in [Mockenhaupt 1996] that the extension operator  $E$  defined by  $\Phi = \xi^t M(t) \xi$  is bounded from  $L^2$  to  $L^{2(d+2k)/d}$ .

In order to obtain estimates for some  $q < 2(d + 2k)/d$  and  $p > 2$ , it seems necessary to consider methods other than the  $TT^*$  argument, which solely relies on the decay estimate for the Fourier transform of the surface measure. For this reason we wish to consider the bilinear restriction estimates for surfaces of codimension greater than 1 and try to obtain the best possible estimates.

Let  $S_1, S_2$  be closed cubes contained in  $I^d$  and define

$$E_i f(x, t) = \int_{S_i} e^{2\pi i(x \cdot \xi + t \cdot \Phi(\xi))} f(\xi) d\xi, \quad i = 1, 2.$$

Let us consider the estimate

$$\|E_1 f E_2 g\|_{L^q(\mathbb{R}^{d+k})} \leq C \|f\|_{L^p(\mathbb{R}^d)} \|g\|_{L^p(\mathbb{R}^d)}. \tag{1-1}$$

For the elliptic surfaces, bilinear estimates can be thought of as a generalization of linear estimates, since a linear restriction estimate follows from the corresponding bilinear one by an argument involving a Whitney decomposition; see, e.g., [Tao et al. 1998]. The advantage of the bilinear estimates is that a wider range of boundedness is possible than for the linear estimate, provided that a separation condition holds between the supports of the functions  $f, g$ . For surfaces with codimension 1, the sharp bilinear (adjoint) restriction estimate for the cone was obtained by Wolff [2001], and for the paraboloid the corresponding estimate was proved by Tao [2003]. The bilinear approach has also been applied to the restriction problem for hyperbolic surfaces: Vargas [2005] used it for the saddle surface in  $\mathbb{R}^3$  and, independently, Lee [2006] proved the bilinear estimate by extending Tao’s method.<sup>1</sup> From these bilinear restriction estimates the corresponding linear ones have been obtained as well.

---

<sup>1</sup>For more general negatively curved surfaces in  $\mathbb{R}^3$  and higher dimensions, Lee [2006] showed the bilinear restriction estimates. However, in higher dimensions the linear estimate could not be deduced from the bilinear one, because the separation condition needed to prove the bilinear estimate for hyperbolic surfaces was more complex than that for the elliptic surfaces.

In order to state our results, we first introduce some notation. For  $v_1, v_2 \in I^d$ , we define the  $k \times d$  matrix  $\mathbf{D}(v_1, v_2)$  by

$$\mathbf{D}(v_1, v_2) = \begin{pmatrix} \nabla\varphi_1(v_2) - \nabla\varphi_1(v_1) \\ \vdots \\ \nabla\varphi_k(v_2) - \nabla\varphi_k(v_1) \end{pmatrix}.$$

Here  $\nabla\varphi_j$  is a row vector. Let  $H\varphi$  denote the Hessian of  $\varphi$  and  $\mathbf{D}^t(v_1, v_2)$  be the transpose of  $\mathbf{D}(v_1, v_2)$ . The following is our main theorem.

**Theorem 1.1.** *Let  $t = (t_1, \dots, t_k)$ ,  $k \geq 1$ . Suppose that, for  $v \in S_1 \cup S_2$  and  $|t| = 1$ ,*

$$\det\left(\sum_{i=1}^k t_i H\varphi_i(v)\right) \neq 0 \tag{1-2}$$

*and, for  $v_1 \in S_1, v_2 \in S_2, |t| = 1$  and for  $v = v_1, v_2$ ,*

$$\det\left[\mathbf{D}(v_1, v_2)\left(\sum_{j=1}^k t_j H\varphi_j(v)\right)^{-1} \mathbf{D}^t(v_1, v_2)\right] \neq 0. \tag{1-3}$$

*Then, for*

$$q > \frac{d + 3k}{d + k} \quad \text{and} \quad \frac{1}{p} + \frac{d + 3k}{d + k} \frac{1}{2q} < 1,$$

*the estimate (1-1) holds.*

As special cases of Theorem 1.1, one can deduce the known bilinear restriction theorems for the elliptic surfaces in [Tao 2003] and the negatively curved ones in [Vargas 2005; Lee 2006].

Let us set

$$\mathbf{M}(t, v_1, v_2, v) := \begin{pmatrix} 0 & \mathbf{D}(v_1, v_2) \\ \mathbf{D}^t(v_1, v_2) & \sum_{i=1}^k t_i H\varphi_i(v) \end{pmatrix}.$$

Assuming the condition (1-2), it is easy to see that (1-3) is equivalent to

$$\det \mathbf{M}(t, v_1, v_2, v) \neq 0 \tag{1-4}$$

for  $v_1 \in S_1, v_2 \in S_2, |t| = 1$  and for  $v = v_1, v_2$ . (One can use the block matrix formula  $\det\begin{pmatrix} A & B \\ C & D \end{pmatrix} = \det(D) \det(A - BD^{-1}C)$ .) The condition (1-4) may seem rather complicated, but such a condition appears naturally when one considers the bilinear  $L^2 \times L^2 \rightarrow L^2$  estimate. When  $k = 1$ , it is closely related to the ‘‘rotational curvature’’; see [Lee 2006] for more details. The necessity of the condition (1-4) will become clear in the course of the proof of Proposition 1.3 below.

From the condition (1-3) it follows that  $\mathbf{D}(v_1, v_2)$  has rank  $k$ . So, the vectors

$$\{\nabla\varphi_i(v_2) - \nabla\varphi_i(v_1) : i = 1, \dots, k\}$$

are linearly independent. This means  $d \geq k$ . If  $d = k$ , then (1-4) implies (1-3), but otherwise (1-4) may hold without (1-3) being satisfied.

In fact, it is possible to obtain a local version (Theorem 1.2 below) of Theorem 1.1, which holds under a weaker assumption. Let  $\mathbf{n}_1, \dots, \mathbf{n}_{d-k}$  be orthonormal vectors (seen as row vectors), which are perpendicular to the span of  $\{\nabla\varphi_i(v_2) - \nabla\varphi_i(v_1) : i = 1, \dots, k\}$  and set

$$N(v_2, v_1) = \begin{pmatrix} \mathbf{n}_1 \\ \vdots \\ \mathbf{n}_{d-k} \end{pmatrix}.$$

Then we can replace the condition (1-4) with

$$\det \left[ N(v_2, v_1) \left( \sum_{i=1}^k t_i H\varphi_i(v) \right) N^t(v_2, v_1) \right] \neq 0 \tag{1-5}$$

whenever  $v_1 \in S_1, v_2 \in S_2, |t| = 1$  and  $v = v_1, v_2$ . It is easy to see that the value of this determinant is independent of the particular choice of orthonormal vectors  $\mathbf{n}_1, \dots, \mathbf{n}_{d-k}$ , and that the condition (1-5) is equivalent to (1-4) under the assumption (1-2).<sup>2</sup> If we have (1-5) instead of (1-3), then we don't need (1-2) to get (1-6) for any  $\alpha > 0$ . More precisely, we have

**Theorem 1.2.** *Suppose that, for any  $v_1 \in S_1, v_2 \in S_2$ , the vectors  $\nabla\varphi_i(v_2) - \nabla\varphi_i(v_1), i = 1, \dots, k$ , are linearly independent and that (1-5) holds for  $v_1 \in S_1, v_2 \in S_2, |t| = 1$  and for  $v = v_1, v_2$ . Then, for any  $\alpha > 0$ , there is a constant  $C_\alpha$  such that*

$$\|E_1 f E_2 g\|_{L^{(d+3k)/(d+k)}(Q_R)} \leq C_\alpha R^\alpha \|f\|_2 \|g\|_2, \tag{1-6}$$

where  $Q_R$  is a cube of side length  $R \gg 1$ .

However, to obtain the global estimates  $L^2 \times L^2 \rightarrow L^q$ , for  $q > (d + 3k)/(d + k)$ , we need to impose a decay condition on the Fourier transform of the surface measure, since it is needed to apply the epsilon removal lemma [Bourgain and Guth 2011]. Under the condition (1-2) such a decay estimate follows from the stationary phase method.

For  $q \geq 2$ , the estimate (1-1) is relatively easier to prove under the conditions (1-2), (1-3). The following may be thought of as a generalization of Theorem 2.3 in [Tao et al. 1998], which is concerned with elliptic hypersurfaces; see also Theorem 4.2 in [Moyua et al. 1999]. A generalization to general hypersurfaces had already been observed in [Lee 2006]. As a byproduct this gives estimates for the endpoint cases of  $(p, q)$  satisfying

$$\frac{1}{p} + \frac{d + 3k}{d + k} \frac{1}{2q} = 1, \quad q \geq 2.$$

**Proposition 1.3.** *Suppose the condition (1-4) holds for  $v_1 \in S_1, v_2 \in S_2$  and  $|t| = 1$ . Then, for  $q \geq 2$  and*

$$\frac{1}{p} + \frac{d + 3k}{d + k} \frac{1}{2q} \leq 1,$$

*the estimate (1-1) holds.*

<sup>2</sup>Indeed, if  $H, N$  and  $D$  are matrices of sizes  $d \times d, (d-k) \times d$  and  $k \times d$ , respectively, such that  $ND^t = 0, \det H \neq 0$  and  $\text{rank}(N^t D^t) = d$ , then  $\det(NHN^t) \neq 0$  if and only if  $\det(DH^{-1}D^t) \neq 0$  because  $\begin{pmatrix} NH \\ D \end{pmatrix} (N^t D^t) = \begin{pmatrix} NHN^t & NHD^t \\ 0 & DD^t \end{pmatrix}$  and  $\begin{pmatrix} N \\ DH^{-1} \end{pmatrix} (N^t D^t) = \begin{pmatrix} NN^t & 0 \\ DH^{-1}N^t & DH^{-1}D^t \end{pmatrix}$ .

**Remark 1.4.** In the proof of the above results we may assume that the aforementioned conditions hold uniformly, by breaking up the extension operator by decomposing  $S_1, S_2$  into sufficiently small pieces. That is to say, there is a constant  $c > 0$  such that for  $v \in S_1 \cup S_2$  and  $|t| = 1$ ,

$$\left| \det \left( \sum_{i=1}^k t_i H \varphi_i(v) \right) \right| \geq c \tag{1-7}$$

and, for  $v_1, v'_1 \in S_1, v_2, v'_2 \in S_2, |t| \sim 1$  and for  $v \in S_1 \cup S_2$ ,

$$\left| \det \left[ \mathbf{D}(v_1, v_2) \left( \sum_{j=1}^k t_j H \varphi_j(v) \right)^{-1} \mathbf{D}^t(v'_1, v'_2) \right] \right| \geq c. \tag{1-8}$$

The same holds also for the conditions (1-4) and (1-5).

**Necessary conditions for (1-1).** By modifying the examples in [Tao and Vargas 2000] with some specific surfaces we see that (1-1) cannot hold in general, unless

$$q \geq \frac{d+k}{d}, \tag{1-9}$$

$$\frac{1}{p} + \frac{d+3k}{d+k} \frac{1}{2q} \leq 1, \tag{1-10}$$

$$\frac{2(d-k)}{p} + \frac{d+3k}{q} \leq 2d. \tag{1-11}$$

In fact,

- (i) (1-9) is necessary for (1-1) to hold under (1-2), and
- (ii) so is (1-10) under the assumption that the matrix  $\mathbf{D}(v_1, v_2)$  has rank  $k$  for  $v_j \in S_j, j = 1, 2$ .

However, in general, (1-11) is not necessarily required for (1-1), but as is well known there are various  $\Phi$  satisfying (1-2) and (1-3) for which (1-1) fails if

$$\frac{2(d-k)}{p} + \frac{d+3k}{q} > 2d.$$

We show (i) and (ii) in the following paragraphs.

(i) By making use of the stationary phase method together with the condition (1-2) it is not difficult to see that, with suitable choice of  $x_0$ , there is a cube  $Q$  of side length  $R \gg 1$  such that  $|E_1(e^{-2\pi i x_0 \cdot \xi} \psi)| \sim |E_2 \psi(x)| \sim R^{-d/2}$  on  $Q$  provided that supports of  $\psi_1, \psi_2$  are small enough. We insert these into (1-1) to see  $R^{-d/2} R^{-d/2} R^{(d+k)/q} \lesssim 1$ , from which we get (1-9) by letting  $R \rightarrow \infty$ . (This can also be shown by making use of a wave packet decomposition, see Lemma 4.2, and randomization.)

(ii) For  $j = 1, 2$ , let  $\Sigma_j$  be the surface  $\{(\xi, \Phi(\xi)) : \xi \in S_j\}$ , and denote by  $d\sigma_j$  the induced Lebesgue measure on  $\Sigma_j$ . To see (1-9) it is more convenient to consider  $f \rightarrow \widehat{f d\sigma_j}$ , instead of dealing with the

operator  $E_j$ . Also, let  $v_j$  be the center of cube  $S_j$  and let  $\zeta_j = (v_j, \Phi(v_j)) \in \Sigma_j$ ,  $j = 1, 2$ . The normal space  $N_j$  to  $\Sigma_j$  at  $\zeta_j$  is spanned by

$$n_{j,i} = (-\nabla\varphi_i(v_j), e_i), \quad i = 1, 2, \dots, k,$$

where  $e_i \in \mathbb{R}^k$  is the usual unit vector with its  $i$ -th entry being equal to 1. Clearly, these vectors are linearly independent because  $D(v_1, v_2)$  has rank  $k$ . Let  $p_n$ ,  $n = 1, \dots, d - k$ , be an orthonormal basis of the orthogonal complement of  $\text{span}\{n_{j,i} : i = 1, 2, \dots, k, j = 1, 2\}$ . Let us set, for  $j = 1, 2$ ,

$$\Lambda_j = \{\zeta \in \Sigma_j : |(\zeta - \zeta_j) \cdot n_{3-j,i}| \leq \delta, |(\zeta - \zeta_j) \cdot p_n| \leq \delta^{\frac{1}{2}}, i = 1, \dots, k, n = 1, \dots, d - k\}.$$

Now, we set  $f_j = \chi_{\Lambda_j}$ ,  $j = 1, 2$ . Then it is easy to see  $|\widehat{f_j d\sigma_j}(x, t)| \gtrsim \delta^{(d+k)/2}$ ,  $j = 1, 2$ , provided that

$$|(x, t) \cdot n_{\ell,i}| \leq c\delta^{-1}, \quad |(x, t) \cdot p_n| \leq c\delta^{-\frac{1}{2}}, \quad i = 1, \dots, k, \ell = 1, 2, n = 1, \dots, d - k$$

with sufficiently small  $c > 0$ . (For example, see the proof Lemma 4.2.) Since (1-1) implies

$$\|\widehat{f_1 d\sigma_1} \widehat{f_2 d\sigma_2}\|_q \lesssim \|f_1\|_p \|f_2\|_p,$$

we get  $\delta^{d+k-(d+3k)/(2q)} \leq C\delta^{(d+k)/p}$  and (1-10) by letting  $\delta \rightarrow 0$ .

**Restriction to complex surfaces.** Using the above theorem we can obtain a bilinear restriction estimate for complex quadratic surfaces. To define the (Fourier) extension operator for a complex surface we first distinguish the dot product and the inner product for complex variables, and define an auxiliary product  $\odot$ . For  $z, w \in \mathbb{C}^m$ , we define  $z \cdot w$ ,  $\langle z, w \rangle$ ,  $z \odot w$  by

$$z \cdot w = \sum_{j=1}^m z_j w_j, \quad \langle z, w \rangle = \sum_{j=1}^m z_j \bar{w}_j, \quad z \odot w = \text{Re}\langle z, w \rangle.$$

Hence, if  $z = x + iy$  and  $w = u + iv$  for  $x, y, u, v \in \mathbb{R}^m$ , then  $z \odot w = x \cdot u + y \cdot v$ . If we identify  $\mathbb{C}^m$  with  $\mathbb{R}^{2m}$  in the usual way, then  $z \odot w$  is just the inner product on  $\mathbb{R}^{2m}$ .

Let  $n \geq 1$  be an integer and let  $D$  be a real symmetric invertible matrix. Then we define the complex quadratic surface  $\gamma \subset \mathbb{C}^{n+1}$  by

$$\gamma(z) = \left(z, \frac{1}{2} z^t D z\right), \quad z \in \mathbb{C}^n. \tag{1-12}$$

Now we define the extension operator  $E_\gamma f$  by

$$E_\gamma f(w) = \int_{\mathbb{C}^n} e^{2\pi i [w \odot \gamma(z)]} f(z) dz, \quad w \in \mathbb{C}^{n+1},$$

where we have written  $dz$  for  $dx dy$ ,  $z = x + iy$ . The operator  $E_\gamma f$  is an extension operator for surfaces of codimension 2 in  $\mathbb{R}^{2n}$ , which is given by  $(x, y, \frac{1}{2} \text{Re}(x + iy)^t D(x + iy), \frac{1}{2} \Im(x + iy)^t D(x + iy))$ ,  $x, y \in \mathbb{R}^n$ . From Theorem 1.1 we can establish the following.

**Corollary 1.5.** *Let  $S_1, S_2$  be closed cubes in  $\mathbb{C}^n$ . Suppose that, for any  $z_1 \in S_1$  and  $z_2 \in S_2$ ,*

$$|(z_2 - z_1)^t D(z_2 - z_1)| \neq 0. \tag{1-13}$$



Then, whenever  $f, g$  are supported on  $S_1, S_2$ , respectively, for

$$q > \frac{n+3}{n+1} \quad \text{and} \quad \frac{1}{p} + \frac{n+3}{n+1} \frac{1}{2q} < 1,$$

there is a constant  $C$  such that

$$\|E_\gamma f \cdot E_\gamma g\|_{L^q(\mathbb{C}^{n+1})} \leq C \|f\|_{L^p(\mathbb{C}^n)} \|g\|_{L^p(\mathbb{C}^n)}.$$

This theorem can also be stated without using the complex number notation, but its use makes it easier to derive the linear estimates from the bilinear one. The condition (1-13) in  $\mathbb{C}^2$  can be contrasted with that in  $\mathbb{R}^2$ . If  $S_1, S_2 \subset \mathbb{R}^2$  and the eigenvalues of  $D$  have the same sign, then the condition (1-13) is always valid if  $\text{dist}(S_1, S_2) \neq 0$ . But, when  $S_1, S_2 \subset \mathbb{C}^2$ , the condition (1-13) may fail even if the separation condition is satisfied. For instance, if  $D$  is the  $2 \times 2$  identity matrix, the condition (1-13) becomes  $|(v_1 - w_1)^2 + (v_2 - w_2)^2| \gtrsim 1$  with  $z_1 = (v_1, v_2)$  and  $z_2 = (w_1, w_2)$ . Since we may factor  $(v_1 - w_1)^2 + (v_2 - w_2)^2$  as  $[(v_1 - w_1) + i(v_2 - w_2)][(v_1 - w_1) - i(v_2 - w_2)]$ , the expression  $|(v_1 - w_1)^2 + (v_2 - w_2)^2|$  may vanish even if  $\text{dist}(S_1, S_2) \gtrsim 1$ . When  $D$  has eigenvalues with different signs, this phenomenon may occur even when  $S_1, S_2 \subset \mathbb{R}^2$ ; for instance, if  $D$  is the  $2 \times 2$  diagonal matrix with diagonal entries 1 and  $-1$ , then we have

$$x \cdot Dx = x_1^2 - x_2^2 = (x_1 + x_2)(x_1 - x_2).$$

This real-variable case was studied by Lee [2006] and Vargas [2005]. In the special case that the surface is two-dimensional they could deduce a linear estimate from the bilinear one.

By adapting their argument, we can obtain the following linear estimate.

**Theorem 1.6.** *Let  $n = 2$  and  $\gamma$  be given by (1-12) with a nonsingular real symmetric matrix  $D$ . Then, for  $q > \frac{10}{3}$  and  $\frac{1}{p} + \frac{2}{q} < 1$ ,*

$$\|E_\gamma f\|_{L^q(\mathbb{C}^3)} \leq C \|f\|_{L^p(\mathbb{C}^2)} \tag{1-14}$$

whenever  $f$  is supported in a bounded set.

By analogy with the corresponding problem for the paraboloid (elliptic or hyperbolic) in  $\mathbb{R}^3$ , it may be conjectured that (1-14) holds if and only if  $q > 3$  and  $\frac{1}{p} + \frac{2}{q} \leq 1$ . Theorem 1.6 extends the known  $(p, q)$  range for the operator  $E_\gamma f$  when  $D$  is a nonsingular real symmetric matrix. This result is an analog of the adjoint Fourier restriction estimates for the hyperbolic paraboloid in  $\mathbb{R}^3$ , which is known to hold for the same range of  $p, q$ . As a special case of the results by Christ [1982, Lemma 4.3] and Mockenhaupt [1996, Theorem 2.11], it was previously known that  $Ef$  maps  $L^2(\mathbb{R}^4)$  boundedly to  $L^4(\mathbb{R}^6)$ . Also, the slightly stronger Lorentz space estimate  $\|Ef\|_{L^{4,2}(\mathbb{R}^6)} \leq C \|f\|_{L^2(\mathbb{R}^4)}$  can be deduced by applying Theorem 1.1 in [Bak and Seeger 2011]. It is quite likely that the multilinear approach will yield further progress on these problems. We hope to return to this problem in the near future.

**Notation.** We adopt the usual convention to let  $C$  or  $c$  represent strictly positive constants, whose value may vary from line to line. But these constants will always be independent of  $f$ , for instance. We write  $A \lesssim B$  or  $B \gtrsim A$  to mean  $A \leq CB$ , and  $A \sim B$  means both  $A \lesssim B$  and  $B \lesssim A$ .

2.  $L^{\frac{4(d+k)}{3d+k}} \times L^{\frac{4(d+k)}{3d+k}} \rightarrow L^2$  estimates and proof of Proposition 1.3

In this section we show Proposition 1.3. Our proof here is different from that in [Tao et al. 1998]. Instead of making use of the boundedness of the averaging operator, we directly exploit the oscillatory decay estimate which is concealed in the averaging operator. For this we need the following lemma.

**Lemma 2.1** [Greenleaf and Seeger 2002, Section 1.1]. *Let  $a \in C_c^\infty(\mathbb{R}^d \times \mathbb{R}^d \times \mathbb{R}^N)$  and set*

$$T_\lambda f(x) = \int_{\mathbb{R}^d} \int_{\mathbb{R}^N} e^{i\lambda\phi(x,y,\theta)} a(x,y,\theta) d\theta f(y) dy,$$

where  $\phi$  is a smooth function on the support of  $a$ . Suppose

$$\det \begin{pmatrix} \phi''_{\theta\theta} & \phi''_{x\theta} \\ \phi''_{y\theta} & \phi''_{xy} \end{pmatrix} \neq 0$$

on the support of  $a$  whenever  $\phi'_\theta = 0$ . Then,  $\|T_\lambda f\|_2 \lesssim \lambda^{-(d+N)/2} \|f\|_2$ .

*Proof of Proposition 1.3.* By interpolation with the trivial  $L^1 \times L^1 \rightarrow L^\infty$  estimate, it suffices to show

$$\|E_1 f_1 E_2 f_2\|_2 \lesssim \|f_1\|_{\frac{4(d+k)}{3d+k}} \|f_2\|_{\frac{4(d+k)}{3d+k}}.$$

For fixed  $\xi_2$ , set

$$\Phi^{\xi_2}(\xi_1, \eta_1) = \Phi(\xi_1) + \Phi(\xi_2) - \Phi(\eta_1) - \Phi(\xi_1 + \xi_2 - \eta_1)$$

and

$$I^{\xi_2}(f_1, \bar{f}_1) = \iint \delta(\Phi^{\xi_2}(\xi_1, \eta_1)) f_1(\xi_1) \bar{f}_1(\eta_1) d\xi_1 d\eta_1,$$

where  $\delta$  is the delta function. Its composition is well defined, since the vectors  $\nabla\phi_i(v_2) - \nabla\phi_i(v_1)$ ,  $i = 1, \dots, k$ , are linearly independent.

By Plancherel's theorem

$$\begin{aligned} \|E_1 f_1 E_2 f_2\|_2^2 &= \iiint \delta(\xi_1 + \xi_2 - \eta_1 - \eta_2, \Phi(\xi_1) + \Phi(\xi_2) - \Phi(\eta_1) - \Phi(\eta_2)) \\ &\quad \times f_1(\xi_1) f_2(\xi_2) \bar{f}_1(\eta_1) \bar{f}_2(\eta_2) d\xi_1 d\xi_2 d\eta_1 d\eta_2 \\ &= \iiint \delta(\Phi^{\xi_2}(\xi_1, \eta_1)) f_1(\xi_1) \bar{f}_1(\eta_1) f_2(\xi_2) \bar{f}_2(\xi_1 + \xi_2 - \eta_1) d\xi_1 d\xi_2 d\eta_1, \end{aligned}$$

where  $f_1, f_2$  are assumed to be supported in  $S_1, S_2$ , respectively. We claim that

$$\|E_1 f_1 E_2 f_2\|_2^2 \lesssim \|f_1\|_{p,1} \|f_2\|_1 \|\bar{f}_1\|_{p,1} \|\bar{f}_2\|_\infty, \tag{2-1}$$

where  $p = (d+k)/d$ . Here  $\|f\|_{r,s}$  denotes the norm of Lorentz space  $L^{r,s}$ . For this we may obviously assume that the functions  $f_1, \bar{f}_1, f_2, \bar{f}_2$  are nonnegative. In order to show (2-1) it suffices to prove

$$|I^{\xi_2}(f, g)| \lesssim \|f\|_{p,1} \|\bar{g}\|_{p,1}. \tag{2-2}$$

Let  $\psi$  be a smooth function with compact Fourier support contained in  $B(0, 1)$  such that  $\hat{\psi} = 1$  on  $B(0, \frac{1}{2})$ . Since  $h(0) = \lim_{j \rightarrow \infty} 2^{jk} \int_{\mathbb{R}^k} \psi(2^j x) h(x) dx$  for any Schwartz function  $h$ , we have

$\delta = \lim_{j \rightarrow \infty} 2^{jk} \psi(2^j x)$ . So, we may write

$$\delta = \sum_{j=-\infty}^{\infty} [2^{(j+1)k} \psi(2^{j+1} x) - 2^{jk} \psi(2^j x)] = \sum_{j=-\infty}^{\infty} 2^{jk} \eta(2^j x),$$

where  $\eta(x) := 2^k \psi(2x) - \psi(x)$ . By the choice of  $\psi$  we see that the Fourier support of  $\eta$  is contained in  $\{\xi : \frac{1}{2} < |\xi| \leq 2\}$ . We decompose  $I^{\xi_2}(f, g)$  by making use of the above decomposition of  $\delta$  to get

$$I^{\xi_2}(f, g) = \sum_{j=-\infty}^{\infty} I_j(f, g),$$

where

$$I_j(f, g) := 2^{kj} \iint \eta(2^j \Phi^{\xi_2}(\xi_1, \eta_1)) f(\xi_1) g(\eta_1) d\xi_1 d\eta_1.$$

It should be noted that we are assuming that  $f, g$  are supported on  $S_1$  and  $\xi_1 + \xi_2 - \eta_1 \in S_2$ . Using the Fourier transform we write  $I_j(f_1, \bar{f}_2)$  as

$$I_j(f, g) = 2^{kj} \int \left( \iint \hat{\eta}(\tau) e^{2^j \tau \cdot \Phi^{\xi_2}(\xi_1, \eta_1)} d\tau f(\xi_1) d\xi_1 \right) g(\eta_1) d\eta_1.$$

Now, we will apply Lemma 2.1 to the double integral inside the parentheses. If we set  $\phi(\xi_1, \eta_1, \tau) = \tau \cdot \Phi^{\xi_2}(\xi_1, \eta_1)$ , then

$$\left| \det \begin{pmatrix} \phi''_{\tau\tau} & \phi''_{\tau\xi_1} \\ \phi''_{\eta_1\tau} & \phi''_{\xi_1\eta_1} \end{pmatrix} \right| = \left| \det \begin{pmatrix} 0 & \mathbf{D}(\xi_1, \xi_1 + \xi_2 - \eta_1) \\ \mathbf{D}(\eta_1, \xi_1 + \xi_2 - \eta_1)^t & \sum_{j=1}^k \tau_j H\varphi_j(\xi_1, \xi_1 + \xi_2 - \eta_1) \end{pmatrix} \right|.$$

So, by the condition (1-4) the last expression does not vanish since  $|\tau| \sim 1$ . Hence, by Lemma 2.1 it follows that

$$|I_j(f, g)| \lesssim 2^{-j \frac{d-k}{2}} \|f\|_2 \|g\|_2.$$

On the other hand, we have the trivial bound  $|I_j(f, g)| \lesssim 2^{kj} \|f\|_1 \|g\|_1$ . Now we may use a summation method (usually called Bourgain’s summation trick) to obtain (2-2).

Considering  $(f_1, \bar{f}_1, f_2, \bar{f}_2) \rightarrow \|E f_1 E f_2\|_2^2$  as a quadrilinear mapping (replacing  $\bar{f}_1, \bar{f}_2$  on the left-hand side by  $\bar{f}_3$  and  $\bar{f}_4$ , respectively), we apply Christ’s multilinear trick [1985]. By symmetry and interpolation we get the estimates

$$\left| \iint E f_1 E f_2 \overline{E f_3 E f_4} dx dt \right| \lesssim \prod_{j=1}^4 \|f_j\|_{p_j, 1}$$

for  $(\frac{1}{p_1}, \frac{1}{p_2}, \frac{1}{p_3}, \frac{1}{p_4})$  contained in the convex hull of the four points

$$v_1 = \left(\frac{1}{p}, \frac{1}{p}, 1, 0\right), \quad v_2 = \left(\frac{1}{p}, \frac{1}{p}, 0, 1\right), \quad v_3 = \left(1, 0, \frac{1}{p}, \frac{1}{p}\right), \quad v_4 = \left(0, 1, \frac{1}{p}, \frac{1}{p}\right)$$

which is contained in the 3-plane  $\Pi = \{u_1 + u_2 + u_3 + u_4 = 1 + \frac{2}{p}\}$ . The convex hull has a nonempty interior in  $\Pi$ , because  $\det(v_1, v_2, v_3, v_4) \neq 0$  as long as  $\frac{1}{p} \neq \frac{1}{2}$ . Hence we may apply the multilinear trick to get

$$\|E f_1 E f_2\|_2^2 \lesssim \|f_1\|_{\frac{4(d+k)}{3d+k}, 4} \|f_2\|_{\frac{4(d+k)}{3d+k}, 4} \|\bar{f}_1\|_{\frac{4(d+k)}{3d+k}, 4} \|\bar{f}_2\|_{\frac{4(d+k)}{3d+k}, 4}. \quad \square$$

### 3. Transversality and the curvature conditions

In this section we prove several lemmas that will play crucial roles in proving Theorem 1.1. These lemmas are related to the curvature conditions.

For  $R \gg 1$  and  $v \in S_1 \cup S_2$ , we set

$$\pi_v = \left\{ (x, t) : \left| x + \left( \sum_{j=1}^k t_j \nabla \varphi_j(v) \right) \right| \leq R^{\frac{1}{2}} \right\}, \quad R^\delta \pi_v = \pi_v + O(R^{\frac{1}{2}+\delta}).$$

Here, for any set  $A \subset \mathbb{R}^{d+k}$  and  $\rho > 0$ , we have  $A + O(\rho) = \{u \in \mathbb{R}^{d+k} : \text{dist}(u, A) \leq C\rho\}$ .

**Lemma 3.1.** *Suppose that the vectors  $\nabla \varphi_j(v_2) - \nabla \varphi_j(v_1)$ ,  $1 \leq j \leq k$ , are linearly independent for all  $v_1 \in S_1$  and  $v_2 \in S_2$ . Then, there is a constant  $C$  such that*

$$\pi_{v_1} \cap \pi_{v_2} \subset B(0, CR^{\frac{1}{2}}).$$

*Proof.* Since the set  $\{\nabla \varphi_j(v_2) - \nabla \varphi_j(v_1)\}_{j=1}^k$  is linearly independent for all  $v_1 \in S_1$  and  $v_2 \in S_2$ , the map  $(t_1, \dots, t_k) \rightarrow (t_1, \dots, t_k)^t \mathbf{D}(v_1, v_2)$  is injective. So, by continuity and compactness it follows that there is a constant  $C$  such that, for all  $v_1 \in S_1$  and  $v_2 \in S_2$ ,

$$|(t_1, \dots, t_k)^t \mathbf{D}(v_1, v_2)| \geq C |(t_1, \dots, t_k)|.$$

If  $(x, t) \in \pi_{v_1} \cap \pi_{v_2}$ , then  $|x + (\sum_{j=1}^k t_j \nabla \varphi_j(v_i))| \leq R^{\frac{1}{2}}$  for  $i = 1, 2$ . This gives

$$|(t_1, \dots, t_k)^t \mathbf{D}(v_1, v_2)| \leq 2R^{\frac{1}{2}}.$$

Hence, the above inequality yields  $|(t_1, \dots, t_k)| \leq CR^{\frac{1}{2}}$ . So, we also get  $|x| \leq CR^{\frac{1}{2}}$ . □

As was already shown in [Lee 2006; Vargas 2005], a simple transversality condition between the two wave packets is not enough to obtain a bilinear estimate beyond the range of the linear  $L^2 \rightarrow L^q$  estimate. So, we need to consider the Fourier supports of the wave packets to put a restriction on the permissible wave packets. This makes the geometry of the associated wave packets more favorable.

For given  $v_1 \in S_1$  and  $v'_2 \in S_2$  we define  $\Pi_1^{v_1, v'_2}$  by

$$\Pi_1^{v_1, v'_2} = \{v'_1 \in S_1 : v'_1 + v'_2 - v_1 \in S_2, \Phi(v_1) + \Phi(v'_1 + v'_2 - v_1) = \Phi(v'_1) + \Phi(v'_2)\}. \quad (3-1)$$

Since  $\{\nabla \varphi_j(v_2) - \nabla \varphi_j(v_1)\}_{j=1}^k$  are linearly independent, by the implicit function theorem we may assume that  $\Pi_1^{v_1, v'_2}$  is a smooth  $(d-k)$ -dimensional surface.<sup>3</sup> We now set


$$\Gamma_1^{v_1, v'_2}(R) = \bigcup_{v'_1 \in \Pi_1^{v_1, v'_2}} R^\delta \pi_{v'_1},$$

which is an  $O(R^{\frac{1}{2}+\delta})$  neighborhood of the conical set with  $k$  null directions. The transversality between  $\Gamma_1^{v_1, v'_2}$  and the opposite plates  $\pi_{v_2}$  is important. Such a transversality is made precise in the following (see Figure 1).

---

<sup>3</sup> We may need to assume that  $S_1$  and  $S_2$  are small enough.





**Figure 1.** Transversality when  $k = 1$  and  $d = 2$ .

**Lemma 3.2.** Let  $0 < \delta \ll 1$ ,  $u \in \mathbb{R}^{d+k}$  and set

$$\tilde{\Gamma}_1^{v_1, v'_2}(R, R^\delta) = \{(x, t) \in \Gamma_1^{v_1, v'_2}(R) : R^{1-\delta} \leq |(x, t)| \leq CR\}.$$

Suppose that the conditions (1-2) and (1-3) hold. Then, if  $S_1$  and  $S_2$  are sufficiently small, there exist a constant  $C$ , independent of  $v_1, v'_2, R$ , and a vector  $u \in \mathbb{R}^{d+k}$  such that for some  $u' \in \mathbb{R}^{d+k}$ ,

$$\tilde{\Gamma}_1^{v_1, v'_2}(R, R^\delta) \cap (R^\delta \pi_{v_2} + u) \subset B(u', CR^{\frac{1}{2}+C\delta}).$$

Note that the set  $\tilde{\Gamma}_1^{v_1, v'_2}(R, R^\delta)$  can be represented as an  $O(R^{\frac{1}{2}+\delta})$  neighborhood of a surface. Let us define the map  $\Phi_1^{v_1, v'_2} : \Pi_1^{v_1, v'_2} \times \mathbb{R}^k \rightarrow \mathbb{R}^{d+k}$  by

$$\Phi_1^{v_1, v'_2}(v, t) = \left( - \sum_{j=1}^k t_j \nabla \varphi_j(v), t \right).$$

Then it is easy to see that

$$\tilde{\Gamma}_1^{v_1, v'_2}(R, R^\delta) \subset \{ \Phi_1^{v_1, v'_2}(v, t) : v \in \Pi_1^{v_1, v'_2}, cR^{1-\delta} \leq |t| \leq CR \} + O(R^{\frac{1}{2}+\delta}).$$

*Proof.* After scaling it is sufficient to show that the intersection of the two sets

$$\Gamma_1 = \{ \Phi_1^{v_1, v'_2}(v, t) : v \in \Pi_1^{v_1, v'_2}, R^{-\delta} \leq |t| \leq C \} + O(R^{-\frac{1}{2}+\delta})$$

and

$$\mathfrak{C}_2(R^{-\frac{1}{2}+\delta}) = \left\{ \left( - \sum_{j=1}^k t_j \nabla \varphi_j(v_2), t \right) : |t| \leq C \right\} + \tilde{u} + O(R^{-\frac{1}{2}+\delta})$$

is contained in a ball of radius  $CR^{-\frac{1}{2}+C\delta}$ . For  $j \geq -C$ , let us set

$$\Gamma_1^j(R^{-\frac{1}{2}+\delta}) = \{ \Phi_1^{v_1, v'_2}(v, t) : v \in \Pi_1^{v_1, v'_2}, 2^{-j-1} \leq |t| \leq 2^{-j} \} + O(R^{-\frac{1}{2}+\delta}).$$

Using homogeneity and a dyadic decomposition in  $t$  for  $\Gamma_1$ , the matter can be reduced to the case  $2^{-1} \leq |t| \leq 1$ . That is to say,

$$\Gamma_1^0(R^{-\frac{1}{2}+\delta}) \cap \mathfrak{C}_2(R^{-\frac{1}{2}+\delta}) \subset B(u, C_0 R^{-\frac{1}{2}+\delta}) \tag{3-2}$$

for some  $u$  and  $C_0 > 0$ . In fact, applying the scaling change of variables  $(x, t) \rightarrow 2^{-j}(x, t)$ , followed by (3-2) and the reverse change of variables, we see that  $\Gamma_1^j(R^{-\frac{1}{2}+\delta}) \cap \mathfrak{C}_2(R^{-\frac{1}{2}+\delta})$  is contained in a

ball of radius  $C_0R^{-\frac{1}{2}+\delta}$ . Since  $\Gamma_1 \subset \bigcup_{2^{-1}R^{-\delta} \leq 2^j \leq C} \Gamma_1^j$ , we know  $\Gamma_1 \cap \mathcal{C}_2(R^{-\frac{1}{2}+\delta})$  is contained in the union of as many as  $\sim \log R$  such balls of radius  $C_0R^{-\frac{1}{2}+\delta}$ . This union of balls is obviously contained in a ball of radius  $CR^{-\frac{1}{2}+C\delta}$  since the set  $\Gamma_1 \cap \mathcal{C}_2(R^{-\frac{1}{2}+\delta})$  is connected.

Since we may assume that  $S_1$  and  $S_2$  are sufficiently small, in order to show (3-2) it is enough to show that the tangent spaces of the surfaces  $\Phi_1^{\nu_1, \nu_2} : \Pi_1^{\nu_1, \nu_2} \times \{2^{-1} \leq |t| \leq 1\} \rightarrow \mathbb{R}^{d+k}$  and  $\{(\sum_{j=1}^k t_j \nabla \varphi_j(\nu_2), t) : |t| \leq C\}$  are uniformly transversal to each other. In fact, since all the underlying sets are compact, by continuity it is enough to check this at each point.

Let  $u_0 = \Phi_1^{\nu_1, \nu_2}(\nu_0, t_0)$  for  $\nu_0 \in \Pi_1^{\nu_1, \nu_2}$  and  $2^{-1} \leq |t_0| \leq 1$ . Let  $\mathbf{v}_1, \dots, \mathbf{v}_{d-k}$  be orthonormal vectors spanning the tangent space  $T_{\nu_0} \Pi_1^{\nu_1, \nu_2}$ . Then the tangent space of the parametrized surface  $\Phi_1^{\nu_1, \nu_2} : \Pi_1^{\nu_1, \nu_2} \times \{2^{-1} \leq |t| \leq 1\} \rightarrow \mathbb{R}^{d+k}$  at  $u_0$  is spanned by the vectors

$$(\nabla \varphi_1(\nu_0), -1, 0, \dots, 0), \quad (\nabla \varphi_2(\nu_0), 0, -1, 0, \dots, 0), \quad \dots, \quad (\nabla \varphi_k(\nu_0), 0, \dots, 0, -1) \tag{3-3}$$

and

$$\left( \mathbf{v}_i \left( \sum_{j=1}^k t_{0,j} H \varphi_j(\nu_0) \right), 0, \dots, 0 \right), \quad i = 1, \dots, d-k. \tag{3-4}$$

On the other hand, the  $k$ -dimensional plane  $\{(-\sum_{j=1}^k t_j \nabla \varphi_j(\nu_2), t) : |t| \leq C\}$  is spanned by

$$(\nabla \varphi_1(\nu_2), -1, 0, \dots, 0), \quad (\nabla \varphi_2(\nu_2), 0, -1, 0, \dots, 0), \quad \dots, \quad (\nabla \varphi_k(\nu_2), 0, \dots, 0, -1). \tag{3-5}$$

Hence it suffices to show that these  $d+k$  vectors are linearly independent, or equivalently that the determinant of the matrix with these vectors as row vectors is nonzero. After Gaussian elimination it is enough to show

$$\det \begin{pmatrix} V(\sum_{j=1}^k t_{0,j} H \varphi_j(\nu_0)) \\ \mathbf{D}(\nu_0, \nu_2) \end{pmatrix} \neq 0, \tag{3-6}$$

where  $V$  is the  $(d-k) \times d$  matrix having  $\mathbf{v}_1, \dots, \mathbf{v}_{d-k}$  as its row vectors. Now by (3-1) we note that the vectors  $\mathbf{v}_1, \dots, \mathbf{v}_{d-k}$  are orthogonal to the span of the vectors

$$\nabla \varphi_j(\nu_0 + \nu_2' - \nu_1) - \nabla \varphi_j(\nu_0), \quad j = 1, \dots, k.$$

Assuming  $S_2$  is small enough, we may replace  $\mathbf{D}(\nu_0, \nu_2)$  by  $\mathbf{D}(\nu_0, \nu_0 + \nu_2' - \nu_1)$ . For simplicity we set  $\tilde{\nu}_2 = \nu_0 + \nu_2' - \nu_1$ . (We may assume there is a  $c > 0$  such that  $|\det[N(\nu_2, \nu_1)(\sum_{i=1}^k t_i H \varphi_i(\nu))N^t(\nu_2, \nu_1)]| > c$  for  $\nu_1 \in S_2$  and  $\nu_2 \in S_2$ ; see Remark 1.4.) Since  $(\sum_{j=1}^k t_{0,j} H \varphi_j(\nu_0))$  is invertible, we need only show

$$\det A \neq 0, \quad \text{where } A = \begin{pmatrix} V \\ \mathbf{D}(\nu_0, \tilde{\nu}_2)(\sum_{j=1}^k t_{0,j} H \varphi_j(\nu_0))^{-1} \end{pmatrix}.$$

Since  $V\mathbf{D}^t(\nu_0, \tilde{\nu}_2) = 0$ , we note that the matrix  $A(V^t \mathbf{D}^t(\nu_0, \tilde{\nu}_2))$  equals

$$\begin{pmatrix} I_{d-k} & 0 \\ \mathbf{D}(\nu_0, \tilde{\nu}_2)(\sum_{j=1}^k t_{0,j} H \varphi_j(\nu_0))^{-1} V^t & \mathbf{D}(\nu_0, \tilde{\nu}_2)(\sum_{j=1}^k t_{0,j} H \varphi_j(\nu_0))^{-1} \mathbf{D}^t(\nu_0, \tilde{\nu}_2) \end{pmatrix}.$$

This matrix is clearly invertible thanks to (1-3). Hence, so is the matrix  $A$ . □

Below we show that the following version of Lemma 3.2 holds, where we assume (1-5) instead of (1-3), dropping the condition (1-2).

**Lemma 3.3.** *Suppose that, for any  $v_1 \in S_1, v_2 \in S_2, \nabla\varphi_i(v_2) - \nabla\varphi_i(v_1), i = 1, \dots, k$  are linearly independent and (1-5) holds for  $v_1 \in S_1, v_2 \in S_2, |t| = 1$  and for  $v = v_1, v_2$ . If  $S_1$  and  $S_2$  are sufficiently small, there is a constant  $C$ , independent of  $v_1, v'_2, R$ , and  $u$  such that, for some  $u' \in \mathbb{R}^{d+1}$ ,*

$$\tilde{\Gamma}_1^{v_1, v'_2}(R, R^\delta) \cap (R^\delta \pi_{v_2} + u) \subset B(u', CR^{\frac{1}{2} + C\delta}).$$

*Proof.* It is sufficient to show that (3-6) holds. As before, under the assumption that  $S_2$  is small enough, we can replace  $\mathbf{D}(v_0, v_2)$  with  $\mathbf{D}(v_0, \tilde{v}_2)$ , where  $\tilde{v}_2 = v_0 + v'_2 - v_1$ . We need only show that

$$\det \begin{pmatrix} V(\sum_{j=1}^k t_{0,j} H\varphi_j(v_0)) \\ \mathbf{D}(v_0, \tilde{v}_2) \end{pmatrix} \neq 0.$$

Since vectors  $v_1, \dots, v_{d-k}$  are orthogonal to the row vectors of  $\mathbf{D}(v_0, \tilde{v}_2)$ , by multiplying the nonsingular matrix  $(V^t \mathbf{D}^t(v_0, \tilde{v}_2))$  by the matrix inside the determinant from the right, we see that the above is equivalent to

$$\det \begin{pmatrix} V(\sum_{j=1}^k t_{0,j} H\varphi_j(v_0)) V^t & V(\sum_{j=1}^k t_{0,j} H\varphi_j(v_0)) \mathbf{D}^t(v_0, \tilde{v}_2) \\ 0 & \mathbf{D}(v_0, \tilde{v}_2) \mathbf{D}^t(v_0, \tilde{v}_2) \end{pmatrix} \neq 0.$$

Since the matrix  $\mathbf{D}(v_0, \tilde{v}_2) \mathbf{D}^t(v_0, \tilde{v}_2)$  is nonsingular, it is clear that the above is equivalent to

$$\det \left[ V \left( \sum_{j=1}^k t_{0,j} H\varphi_j(v_0) \right) V^t \right] \neq 0,$$

which is the condition (1-5). □

#### 4. Proof of Theorem 1.1

In this section we will prove Theorem 1.1. Our proof is similar to that in [Lee 2006]; also see [Tao 2003]. To prove Theorem 1.1, we need only show that, for  $p > (d + 3k)/(d + k)$ ,

$$\|E_1 f E_2 g\|_p \leq C \|f\|_2 \|g\|_2$$

since we can obtain the desired conclusion by interpolating this estimate with the trivial estimate  $\|E_1 f E_2 g\|_\infty \leq \|f\|_1 \|g\|_1$ . By an  $\epsilon$ -removal argument [Tao and Vargas 2000; Bourgain and Guth 2011], it is sufficient to show that (1-6) holds for any  $\alpha > 0$ . In fact, by the assumption that  $\sum_{j=1}^k t_j H\varphi_j(v)$  is nonsingular for  $v \in \text{supp } f \cup \text{supp } g$  as long as  $|t| = 1$ , it follows that

$$|E_\kappa(a_\kappa)(x, t)| \lesssim (|x| + |t|)^{-\frac{d}{2}}, \quad \kappa = 1, 2,$$

where  $a_1, a_2$  are smooth bump functions which vanish on the supports of  $f$  and  $g$ , respectively. This can be shown by the stationary phase method. Hence, the arguments in the papers mentioned above work here without modification.

**Proposition 4.1.** *Let  $0 < \delta \ll 1$ . If (1-6) holds, then for any  $\epsilon > 0$*

$$\|E_1 f E_2 g\|_{L^{(d+3k)/(d+k)}(Q_R)} \leq C_\epsilon R^{\max(\alpha(1-\delta), C\delta)+\epsilon} \|f\|_2 \|g\|_2, \tag{4-1}$$

with  $C$  independent of  $\delta$ .

By iterating finitely many times the implication in Proposition 4.1, we can easily obtain the estimate (1-6) for any  $\alpha > 0$ .

**Wave packet decomposition.** In this section we decompose the function  $Ef$  into wave packets. Let  $R \gg 1$ . We define

$$\mathcal{L} = \mathcal{L}(R) := R^{\frac{1}{2}}\mathbb{Z}^d, \quad \mathcal{V} = \mathcal{V}(R) := R^{-\frac{1}{2}}\mathbb{Z}^d.$$

Let  $\psi$  be a nonnegative Schwartz function such that  $\hat{\psi}$  is supported on  $B(0, 1)$  and  $\sum_{k \in \mathbb{Z}^d} \psi(\cdot - k) = 1$ . Also, let  $\zeta$  be a smooth function supported on  $B(0, 1)$  and  $\sum_{k \in \mathbb{Z}^d} \zeta(\cdot - k) = 1$ .

For  $\ell \in \mathcal{L}$ ,  $\nu \in \mathcal{V}$  we set  $\psi_\ell(x) := \psi((x - \ell)/R^{\frac{1}{2}})$ ,  $\zeta_\nu(\xi) := \zeta(R^{\frac{1}{2}}(\xi - \nu))$ , and for a given function  $f$ , we define  $f_{\ell, \nu}$  by

$$f_{\ell, \nu} = \mathcal{F}(\psi_\ell \mathcal{F}^{-1}(\zeta_\nu f)),$$

where  $\mathcal{F}$ ,  $\mathcal{F}^{-1}$  denote the Fourier transform and the inverse Fourier transform, respectively. Then, it follows that  $f = \sum_{\nu \in \mathcal{V}} \sum_{\ell \in \mathcal{L}} f_{\ell, \nu}$ . Hence we may write

$$Ef = \sum_{\nu \in \mathcal{V}} \sum_{\ell \in \mathcal{L}} Ef_{\ell, \nu}. \tag{4-2}$$

**Lemma 4.2.** *If  $|t| \lesssim R$ , then*

$$|Ef_{\ell, \nu}(x, t)| \leq C_N \left(1 + R^{-\frac{1}{2}} \left|x - \ell + \sum_{j=1}^k t_j \nabla \varphi_j(\nu)\right|\right)^{-N} M(\mathcal{F}^{-1}(\zeta_\nu f))(\ell) \tag{4-3}$$

for all  $N \geq 0$ . Here,  $Mf$  is the Hardy–Littlewood maximal function of  $f$ .

*Proof.* Since  $f_{\ell, \nu}$  is supported in  $B(\nu, 3R^{-\frac{1}{2}})$ , multiplying by a harmless smooth bump function  $\tilde{\chi}$  supported in  $B(0, 5)$  and satisfying  $\tilde{\chi} = 1$  on  $B(0, 3)$ , we may write

$$Ef_{\ell, \nu}(x, t) = \int K(x - z, t) \psi_\ell(z) \mathcal{F}^{-1} f_\nu(z) dz,$$

where  $K(x, t) = \int e^{2\pi i(x \cdot \xi + t \cdot \Phi(\xi))} \chi(R^{\frac{1}{2}}(\xi - \nu)) d\xi$ . Changing variables  $\xi \rightarrow R^{-\frac{1}{2}}\xi + \nu$ ,

$$K(x, t) = R^{-\frac{d}{2}} e^{2\pi i x \cdot \nu} \int e^{2\pi i(R^{-1/2}x \cdot \xi + t \cdot \Phi(R^{-1/2}\xi + \nu))} \chi(\xi) d\xi.$$

Since  $|t| \lesssim R$ , we know  $\nabla_\xi(R^{-\frac{1}{2}}x \cdot \xi + t \cdot \Phi(R^{-\frac{1}{2}}\xi + \nu)) = R^{-\frac{1}{2}}(x + \sum_{j=1}^k t_j \nabla \varphi_j(\nu)) + O(1)$ . This follows by Taylor’s expansion. Hence, by repeated integration by parts we get

$$|K(x, t)| \leq C_N R^{-\frac{d}{2}} \left(1 + R^{-\frac{1}{2}} \left|x + \sum_{j=1}^k t_j \nabla \varphi_j(\nu)\right|\right)^{-N}.$$

Once this is established, (4-3) follows by a standard argument. See [Lee 2006] for the details. □



From the above lemma we see that  $Ef_{\ell,v}$  is essentially supported on

$$\pi_{\ell,v} = \pi_v + (\ell, 0). \tag{4-4}$$

If  $\pi = \pi_{\ell,v}$ , we define  $\nu(\pi) = \nu$ , which may be considered as the (generalized) direction of  $\pi$ .

The following is the main lemma of this section.

**Lemma 4.3.** *Let  $R \gg 1$ . Then,  $Ef$  can be rewritten as*

$$Ef(x, t) = \sum_{(\ell,v) \in \mathcal{L} \times \mathcal{V}} c_{\ell,v} P_{\ell,v}(x, t) \tag{4-5}$$

and  $c_{\ell,v}, P_{\ell,v}$  satisfy the following:

- (i)  $\mathcal{F}(P_{\ell,v}(\cdot, t))$  is supported in the disc  $D(v, CR^{-\frac{1}{2}})$ .
- (ii) If  $|t| \lesssim R$ , then for any  $N \geq 0$

$$|P_{\ell,v}(x, t)| \leq C_N R^{-\frac{d}{4}} \left( 1 + R^{-\frac{1}{2}} \left| x - \ell + \left( \sum_{j=1}^k t_j \nabla \varphi_j(v) \right) \right| \right)^{-N}.$$

- (iii)  $(\sum_{(\ell,v) \in \mathcal{L} \times \mathcal{V}} |c_{\ell,v}|^2)^{\frac{1}{2}} \lesssim \|f\|_2$ .

- (iv) If  $|t| \lesssim R$ , then  $\|\sum_{(\ell,v) \in \mathcal{W}} P_{\ell,v}(\cdot, t)\|_2^2 \lesssim \#\mathcal{W}$  for any  $\mathcal{W} \subset \mathcal{L} \times \mathcal{V}$ .

*Proof.* We define  $c_{\ell,v}$  and  $P_{\ell,v}$  by

$$c_{\ell,v} = R^{\frac{d}{4}} M(\mathcal{F}^{-1} f_\nu)(\ell), \quad P_{\ell,v}(x, t) = c_{\ell,v}^{-1} Ef_{\ell,v}(x, t),$$

where  $M$  denotes the Hardy–Littlewood maximal function. Then we have (4-5) from (4-2). Since  $Ef_{\ell,v}(\cdot, y) = \mathcal{F}^{-1}(e^{2\pi iy \Phi} f_{\ell,v})$ , we know  $Ef_{\ell,v}(\cdot, y)$  has Fourier support contained in  $\text{supp } f_{\ell,v}$ , which is in turn contained in  $D(v, CR^{-\frac{1}{2}})$ . Thus (i) follows and so does (ii) from Lemma 4.2.

In order to show (iii), note that

$$\sum_{(\ell,v) \in \mathcal{L} \times \mathcal{V}} |c_{\ell,v}|^2 = R^{\frac{d}{2}} \sum_{(\ell,v) \in \mathcal{L} \times \mathcal{V}} M(\mathcal{F}^{-1}(\zeta_\nu f))(\ell)^2. \tag{4-6}$$

Since  $\zeta_\nu f$  is supported on  $B(v, CR^{\frac{1}{2}})$ , we have  $M(\mathcal{F}^{-1}(\zeta_\nu f))(x) \sim M(\mathcal{F}^{-1}(\zeta_\nu f))(x')$  if  $|x - x'| \lesssim R^{\frac{1}{2}}$ . Hence, from the Hardy–Littlewood maximal theorem and Plancherel’s theorem we have that, for each  $\nu$ ,

$$R^{\frac{d}{2}} \sum_{(\ell,v) \in \mathcal{L} \times \mathcal{V}} |M(\mathcal{F}^{-1}(\zeta_\nu f))(\ell)|^2 \lesssim \int |M(\mathcal{F}^{-1}(\zeta_\nu f))(x)|^2 dx \lesssim \|\zeta_\nu f\|_2^2.$$

Combining this and (4-6) we obtain  $\sum_{(\ell,v) \in \mathcal{L} \times \mathcal{V}} |c_{\ell,v}|^2 \lesssim \sum_{\nu \in \mathcal{V}} \|\zeta_\nu f\|_2^2 \lesssim \|f\|_2^2$ , and (iii).

Finally, we consider (iv). Since  $\sum_{\ell: (\ell,v) \in \mathcal{W}} P_{\ell,v}(\cdot, t)$  is Fourier-supported in  $D(v, CR^{-\frac{1}{2}})$ , which has bounded overlap as  $\nu$  varies over  $\mathcal{V}$ , by Plancherel’s theorem,

$$\left\| \sum_{(\ell,v) \in \mathcal{W}} P_{\ell,v}(\cdot, t) \right\|_2^2 \lesssim \sum_{\nu \in \mathcal{V}} \left\| \sum_{\ell: (\ell,v) \in \mathcal{W}} P_{\ell,v}(\cdot, t) \right\|_2^2.$$

From (ii) it is easy to see that  $\left\| \sum_{\ell: (\ell, \nu) \in \mathcal{W}} P_{\ell, \nu}(\cdot, t) \right\|_2^2 \lesssim \#\{\ell : (\ell, \nu) \in \mathcal{W}\}$ . Hence, combining this with the above gives (iv).  $\square$

**Dyadic pigeonholing and reduction.** In the remainder of this section we will prove Proposition 4.1. For simplicity we set

$$p_0 = \frac{d + 3k}{d + k}.$$

By translation invariance we may assume that  $Q_R$  is centered at the origin. Let

$$\mathcal{W}_i \subset \{(\ell, \nu) \in \mathcal{L} \times \mathcal{V} : \nu \in S_i + O(R^{-\frac{1}{2}})\}, \quad i = 1, 2.$$

By Lemma 4.3 and the standard reduction with pigeonholing, which may only cause a loss of  $(\log R)^C$ , see [Lee 2006; Tao 2003], the matter is reduced to showing

$$\left\| \sum_{\omega_1 \in \mathcal{W}_1} P_{\omega_1} \sum_{\omega_2 \in \mathcal{W}_2} P_{\omega_2} \right\|_{L^{p_0}(Q_R)} \lesssim (R^{(1-\delta)\alpha} + R^{C\delta})(\#\mathcal{W}_1 \#\mathcal{W}_2)^{\frac{1}{2}}$$

whenever  $P_{\omega_1}, P_{\omega_2}$  satisfy (i), (ii), (iv) in Lemma 4.3. Here  $A \lesssim B$  means  $A \leq C_\epsilon R^\epsilon B$  for any  $\epsilon > 0$ .

By a further pigeonholing argument we specify the associated quantities in dyadic scales. Let  $\mathcal{Q}$  be a collection of almost disjoint cubes of the same side length  $\sim R^{\frac{1}{2}}$  which cover  $Q_R$ . For each  $q \in \mathcal{Q}$  we define

$$\mathcal{W}_j(q) = \{\omega_j \in \mathcal{W}_j : \pi_{\omega_j} \cap R^\delta q \neq \emptyset\}.$$

For dyadic numbers  $\rho_1, \rho_2$  with  $1 \leq \rho_1, \rho_2 \leq R^{100d}$ , we define

$$\mathcal{Q}(\rho_1, \rho_2) = \{q \in \mathcal{Q} : \rho_j \leq \#\mathcal{W}_j(q) < 2\rho_j, \quad j = 1, 2\}. \tag{4-7}$$

For  $\omega \in \mathcal{W}_1 \cup \mathcal{W}_2$ , we set

$$\lambda(\omega; \rho_1, \rho_2) = \#\{q \in \mathcal{Q}(\rho_1, \rho_2) : \pi_\omega \cap R^\delta q \neq \emptyset\}.$$

For a dyadic number  $1 \leq \lambda \leq R^{100d}$ , we define

$$\mathcal{W}_j[\lambda; \rho_1, \rho_2] = \{\omega_j \in \mathcal{W}_j : \lambda \leq \lambda(\omega_j; \rho_1, \rho_2) < 2\lambda\}, \quad j = 1, 2. \tag{4-8}$$

By a standard pigeonhole argument, it is sufficient to show

$$\left( \sum_{q \in \mathcal{Q}(\rho_1, \rho_2)} \left\| \sum_{\omega_1 \in \mathcal{W}_1[\lambda_1; \rho_1, \rho_2]} P_{\omega_1} \sum_{\omega_2 \in \mathcal{W}_2[\lambda_2; \rho_1, \rho_2]} P_{\omega_2} \right\|_{L^{p_0}(q)}^{p_0} \right)^{\frac{1}{p_0}} \lesssim (R^{(1-\delta)\alpha} + R^{C\delta})(\#\mathcal{W}_1 \#\mathcal{W}_2)^{\frac{1}{2}}. \tag{4-9}$$

For the rest of the proof we assume that  $q \in \mathcal{Q}(\rho_1, \rho_2)$ ,  $\omega_1 \in \mathcal{W}_1[\lambda_1; \rho_1, \rho_2]$  and  $\omega_2 \in \mathcal{W}_2[\lambda_2; \rho_1, \rho_2]$  if it is not mentioned otherwise. So, the above sums are denoted simply by  $\sum_q, \sum_{\omega_1}$ , and  $\sum_{\omega_2}$ , respectively.

**Induction argument.** For brevity let us put

$$\Delta = \bigcup_{q \in \mathcal{Q}(\rho_1, \rho_2)} q.$$

Let  $\{B\}$  be a collection of almost disjoint cubes of the same side length  $R^{1-\delta}$ , which cover  $Q_R$ . Then

$$\text{the left-hand side of (4-9)} \leq \sum_B \left\| \sum_{\omega_1} P_{\omega_1} \sum_{\omega_2} P_{\omega_2} \right\|_{L^{p_0}(\Delta \cap B)}. \tag{4-10}$$

We define a relation  $\sim$  between  $\omega_1$  (or  $\omega_2$ ) and the cubes in  $\{B\}$ . For each  $\omega \in \mathcal{W}_1[\lambda_1; \rho_1, \rho_2] \cup \mathcal{W}_2[\lambda_2; \rho_1, \rho_2]$ , we define  $B^*(\omega) \in \{B\}$  to be the cube which maximizes the quantity

$$\#\{q \in \mathcal{Q}(\rho_1, \rho_2) : \pi_\omega \cap R^\delta q \neq \emptyset, q \cap B \neq \emptyset\}. \tag{4-11}$$

Then the relation  $\sim$  is defined as

$$\omega \sim B \quad \text{if } B \cap 10B^*(\omega) \neq \emptyset.$$

Here  $10B^*(\omega)$  is the cube which has the same center as  $B^*(\omega)$  and side length 10 times as large as that of  $B^*(\omega)$ . Using this relation we divide the sum into three parts to get

$$\begin{aligned} \sum_B \left\| \sum_{\omega_1} P_{\omega_1} \sum_{\omega_2} P_{\omega_2} \right\|_{L^{p_0}(\Delta \cap B)} &\leq \sum_B \left\| \sum_{\substack{\omega_1: \\ \omega_1 \sim B}} P_{\omega_1} \sum_{\substack{\omega_2: \\ \omega_2 \sim B}} P_{\omega_2} \right\|_{L^{p_0}(\Delta \cap B)} \\ &\quad + \sum_B \left\| \sum_{\substack{\omega_1: \\ \omega_1 \not\sim B}} P_{\omega_1} \sum_{\substack{\omega_2: \\ \omega_2 \sim B}} P_{\omega_2} \right\|_{L^{p_0}(\Delta \cap B)} + \sum_B \left\| \sum_{\substack{\omega_1: \\ \omega_1 \not\sim B}} P_{\omega_1} \sum_{\substack{\omega_2: \\ \omega_2 \not\sim B}} P_{\omega_2} \right\|_{L^{p_0}(\Delta \cap B)}. \end{aligned}$$

We will first show that

$$\sum_B \left\| \sum_{\substack{\omega_1: \\ \omega_1 \sim B}} P_{\omega_1} \sum_{\substack{\omega_2: \\ \omega_2 \sim B}} P_{\omega_2} \right\|_{L^{p_0}(\Delta \cap B)} \lesssim R^{(1-\delta)\alpha} (\#\mathcal{W}_1 \#\mathcal{W}_2)^{\frac{1}{2}}. \tag{4-12}$$

By applying the hypothesis (1-6), (iv) in Lemma 4.3, and the Cauchy–Schwarz inequality,

$$\sum_B \left\| \sum_{\substack{\omega_1: \\ \omega_1 \sim B}} P_{\omega_1} \sum_{\substack{\omega_2: \\ \omega_2 \sim B}} P_{\omega_2} \right\|_{L^{p_0}(\Delta \cap B)} \leq CR^{(1-\delta)\alpha} \prod_{j=1}^2 \left( \sum_B \#\{\omega_j : \omega_j \sim B\} \right)^{\frac{1}{2}}.$$

From the definition of the relation  $\sim$  it is clear that  $\#\{B : \omega_j \sim B\} \leq C$ . Hence, for  $j = 1, 2$

$$\sum_B \#\{\omega_j : \omega_j \sim B\} = \sum_{\omega_j} \#\{B : \omega_j \sim B\} \lesssim W_j.$$

By inserting this into the previous inequality, we get (4-12).

Now, to prove (4-9) it is enough to show

$$\left\| \sum_{\substack{\omega_1: \\ \omega_1 \sim B}} P_{\omega_1} \sum_{\substack{\omega_2: \\ \omega_2 \not\sim B}} P_{\omega_2} \right\|_{L^{p_0}(\Delta \cap B)} \lesssim R^{C\delta} (\#\mathcal{W}_1 \#\mathcal{W}_2)^{\frac{1}{2}}$$

and

$$\left\| \sum_{\omega_1 \not\sim B} P_{\omega_1} \sum_{\omega_2} P_{\omega_2} \right\|_{L^{p_0}(\Delta \cap B)} \lesssim R^{C\delta} (\#\mathcal{W}_1 \#\mathcal{W}_2)^{\frac{1}{2}}. \tag{4-13}$$

The proofs of these two estimates are similar. So, we will only prove (4-13). By Plancherel’s theorem,  $\|Ef(\cdot, t)\|_2 \leq \|f\|_2$  for all  $t \in \mathbb{R}^k$ . Integration in  $t$  gives  $\|Ef\|_{L^2(Q_R)} \lesssim R^k/2\|f\|_2$ . By the Schwarz inequality it follows that

$$\|E_1 f E_2 g\|_{L^1(Q_R)} \lesssim R^k \|f\|_2 \|g\|_2.$$

Combining this with (iv) in Lemma 4.3 yields

$$\left\| \sum_{\omega_1 \not\sim B} P_{\omega_1} \sum_{\omega_2} P_{\omega_2} \right\|_{L^1(\Delta \cap B)} \lesssim R^k (\#\mathcal{W}_1 \#\mathcal{W}_2)^{\frac{1}{2}}. \tag{4-14}$$

Hence, the (4-13) follows from interpolation between (4-14) and

$$\left\| \sum_{\substack{\omega_1: \\ \omega_1 \not\sim B}} P_{\omega_1} \sum_{\omega_2} P_{\omega_2} \right\|_{L^2(\Delta \cap B)} \lesssim R^{C\delta} R^{-\frac{d-k}{4}} (\#\mathcal{W}_1 \#\mathcal{W}_2)^{\frac{1}{2}}. \tag{4-15}$$

Now it remains to show the  $L^2$ -estimate (4-15).

**$L^2$  estimate.** To prove (4-15) it suffices to show

$$\sum_{\substack{q \in \mathcal{Q}(\rho_1, \rho_2) \\ q \subset 2B}} \left\| \sum_{\substack{\omega_1: \\ \omega_1 \not\sim B}} P_{\omega_1} \sum_{\omega_2} P_{\omega_2} \right\|_{L^2(q)}^2 \lesssim R^{C\delta} R^{-\frac{d-k}{2}} \#\mathcal{W}_1 \#\mathcal{W}_2. \tag{4-16}$$

For  $j = 1, 2$ , let us set

$$\mathcal{W}_j(q) = \{\omega_j \in \mathcal{W}_j[\lambda_j; \rho_1, \rho_2] : \omega_j \cap R^\delta q \neq \emptyset\}, \quad \mathcal{W}_j^{\not\sim B}(q) = \{\omega_j \in \mathcal{W}_j(q) : \omega_j \not\sim B\}.$$

Then by (ii) in Lemma 4.3 we may discard some harmless terms, whose contributions are  $O(R^{-C\delta})$ . Hence, it suffices to show

$$\sum_{\substack{q \in \mathcal{Q}(\rho_1, \rho_2) \\ q \subset 2B}} \left\| \sum_{\omega_1 \in \mathcal{W}_1^{\not\sim B}(q)} P_{\omega_1} \sum_{\omega_2(q)} P_{\omega_2} \right\|_2^2 \lesssim R^{C\delta} R^{-\frac{d-k}{2}} \#\mathcal{W}_1 \#\mathcal{W}_2. \tag{4-17}$$

By using Plancherel’s theorem we write

$$\left\| \sum_{\omega_1 \in \mathcal{W}_1^{\not\sim B}(q)} P_{\omega_1} \sum_{\omega_2 \in \mathcal{W}_2(q)} P_{\omega_2} \right\|_2^2 = \sum_{\omega_1 \in \mathcal{W}_1^{\not\sim B}(q)} \sum_{\omega'_2 \in \mathcal{W}_2(q)} \sum_{\omega'_1 \in \mathcal{W}_1^{\not\sim B}(q)} \sum_{\omega_2 \in \mathcal{W}_2(q)} \langle \hat{P}_{\omega_1} * \hat{P}_{\omega_2}, \hat{P}_{\omega'_1} * \hat{P}_{\omega'_2} \rangle.$$



Let us write  $\omega_j = (\ell_j, \nu_j)$ ,  $\omega'_j = (\ell'_j, \nu'_j)$ ,  $j = 1, 2$ . For any  $\nu_1 \in S_1$ ,  $\nu'_2 \in S_2$ , we define  $\mathcal{W}_1^{\mathcal{R}^B}(q; \nu_1, \nu'_2)$  by

$$\mathcal{W}_1^{\mathcal{R}^B}(q; \nu_1, \nu'_2) = \{\omega'_1 = (\ell'_1, \nu'_1) \in \mathcal{W}_1^{\mathcal{R}^B}(q) : \nu'_1 \in \Pi_1^{\nu_1, \nu'_2} + O(R^{-\frac{1}{2}})\}.$$

Then  $\widehat{P}_{\omega_1} * \widehat{P}_{\omega_2}$  is supported on the  $O(R^{-\frac{1}{2}})$ -neighborhood of the point  $(\nu_1 + \nu_2, \Phi(\nu_1) + \Phi(\nu_2))$ . So the inner product  $\langle \widehat{P}_{\omega_1} * \widehat{P}_{\omega_2}, \widehat{P}_{\omega'_1} * \widehat{P}_{\omega'_2} \rangle$  vanishes unless

$$\nu_1 + \nu_2 = \nu'_1 + \nu'_2 + O(R^{-\frac{1}{2}}), \quad \Phi(\nu_1) + \Phi(\nu_2) = \Phi(\nu'_1) + \Phi(\nu'_2) + O(R^{-\frac{1}{2}}).$$

Thus, for given  $\nu_1$  and  $\nu'_2$ , we see that  $\nu'_1$  is contained in an  $O(R^{-\frac{1}{2}})$ -neighborhood of  $\Pi_1^{\nu_1, \nu'_2}$ , which is defined by (3-1). Once  $\nu_1$ ,  $\nu'_1$  and  $\nu'_2$  are given, then there are only  $O(1)$  many  $\nu_2$ , since  $\nu_2$  should be in an  $O(R^{-\frac{1}{2}})$ -neighborhood of the point  $\nu_1 + \nu'_1 - \nu'_2$ . Therefore,

$$\left\| \sum_{\omega_1 \in \mathcal{W}_1^{\mathcal{R}^B}(q)} P_{\omega_1} \sum_{\omega_2 \in \mathcal{W}_2(q)} P_{\omega_2} \right\|_2^2 \lesssim R^{-\frac{d-k}{2}} \sum_{\omega_1 \in \mathcal{W}_1^{\mathcal{R}^B}(q)} \sum_{\omega'_2 \in \mathcal{W}_2(q)} \#\mathcal{W}_1^{\mathcal{R}^B}(q; \nu_1, \nu'_2),$$

where we also used

$$|\langle P_{\omega_1} P_{\omega_2}, P_{\omega'_1} P_{\omega'_2} \rangle| \lesssim R^{-\frac{d-k}{2}}.$$

This follows from (ii) in Lemma 4.3 and the transversality between  $\pi_{\omega_1}$  ( $\pi_{\omega'_1}$ ) and  $\pi_{\omega_2}$  ( $\pi_{\omega'_2}$ ), respectively. Hence, (4-17) follows if we show

$$\max_{q \subset 2B, \nu_1, \nu'_2} \#\mathcal{W}_1^{\mathcal{R}^B}(q; \nu_1, \nu'_2) \sum_{\substack{q \in \mathcal{Q}(\rho_1, \rho_2) \\ q \subset 2B}} \#\mathcal{W}_1^{\mathcal{R}^B}(q) \#\mathcal{W}_2(q) \lesssim R^{C\delta} \#\mathcal{W}_1 \#\mathcal{W}_2. \tag{4-18}$$

We will prove (4-18), assuming for the moment that

$$\max_{q \subset 2B, \nu_1, \nu'_2} \#\mathcal{W}_1^{\mathcal{R}^B}(q; \nu_1, \nu'_2) \lesssim R^{C\delta} \frac{\#\mathcal{W}_2}{\lambda_1 \rho_2}. \tag{4-19}$$

To this end it is enough to show

$$\sum_{\substack{q \in \mathcal{Q}(\rho_1, \rho_2) \\ q \subset 2B}} \#\mathcal{W}_1^{\mathcal{R}^B}(q) \#\mathcal{W}_2(q) \lesssim \lambda_1 \rho_2 \#\mathcal{W}_1.$$

Recalling  $\#\mathcal{W}_2(q) \lesssim \rho_2$ , we see that the left-hand side is bounded by

$$C\rho_2 \sum_{q \in \mathcal{Q}(\rho_1, \rho_2)} \#\mathcal{W}_1(q).$$

Changing the order of summation, we see this in turn is bounded by  $C\rho_2 \sum_{\omega_1} \#\{q \in \mathcal{Q}(\rho_1, \rho_2) : \pi_{\omega_1} \cap R^\delta q\}$ . Since

$$\#\{q \in \mathcal{Q}(\rho_1, \rho_2) : \pi_{\omega_1} \cap R^\delta q\} \lesssim \lambda_1,$$

the desired inequality (4-18) follows.

**Proof of (4-19).** Fix  $q \subset 2B$ ,  $v_1 \in S_1$  and  $v'_2 \in S_2$ . Let us consider the set

$$\mathbf{S} := \{(\tilde{q}, \omega_1, \omega_2) \in \mathcal{Q}(\rho_1, \rho_2) \times \mathcal{W}_1^{\times B}(q; v_1, v'_2) \times \mathcal{W}_2 : \pi_{\omega_1} \cap R^\delta \tilde{q} \neq \emptyset, \pi_{\omega_2} \cap R^\delta \tilde{q} \neq \emptyset, \text{dist}(\tilde{q}, q) \geq R^{1-\delta}\}.$$

To prove (4-19) it suffices to show

$$R^{-C\delta} \lambda_1 \rho_2 \#\mathcal{W}_1^{\times B}(q; v_1, v'_2) \lesssim \#\mathbf{S} \lesssim R^{C\delta} \#\mathcal{W}_2. \tag{4-20}$$

For the lower bound it is enough to show that, for each  $\omega_1 \in \mathcal{W}_1^{\times B}(q; v_1, v'_2)$ ,

$$\#\{(\tilde{q}, \omega_2) \in \mathcal{Q}(\rho_1, \rho_2) \times \mathcal{W}_2 : \pi_{\omega_1} \cap R^\delta \tilde{q} \neq \emptyset, \pi_{\omega_2} \cap R^\delta \tilde{q} \neq \emptyset, \text{dist}(\tilde{q}, q) \geq R^{1-\delta}\} \geq R^{-C\delta} \lambda_1 \rho_2.$$

By (4-8),  $\omega_1$  contains as many as  $O(\lambda_1)$  cubes  $\tilde{q}$  in  $\mathcal{Q}(\rho_1, \rho_2)$ . (Recall that we are assume assuming  $q \in \mathcal{Q}(\rho_1, \rho_2)$ ,  $\omega_1 \in \mathcal{W}_1[\lambda_1; \rho_1, \rho_2]$  and  $\omega_2 \in \mathcal{W}_2[\lambda_1; \rho_1, \rho_2]$ .) Let  $B^*(\omega_1) \in \mathcal{Q}$  be the cube which maximizes the quantity given by (4-11) with  $\omega = \omega_1$ . Since  $\omega_1 \sim B$ , it follows from the definition of the relation  $\sim$  that  $\text{dist}(B^*(\omega_1), B) \gtrsim R^{1-\delta}$ . Since  $\pi_{\omega_1} + O(R^{\frac{1}{2}+\delta})$  can be covered by  $R^{C\delta}$  cubes  $B$ , by a simple pigeonholing argument we get

$$\#\{\tilde{q} \in \mathcal{Q}(\rho_1, \rho_2) : \pi_{\omega_1} \cap R^\delta \tilde{q} \neq \emptyset, \text{dist}(\tilde{q}, q) \geq R^{1-\delta}\} \gtrsim R^{-C\delta} \lambda_1.$$

Next, for the upper bound it suffices to show that, for any  $\omega_2 \in \mathcal{W}_2$ ,

$$\#\{(\tilde{q}, \omega_1) \in \mathcal{Q}(\rho_1, \rho_2) \times \mathcal{W}_1^{\times B}(q; v_1, v'_2) : \pi_{\omega_1} \cap R^\delta \tilde{q} \neq \emptyset, \pi_{\omega_2} \cap R^\delta \tilde{q} \neq \emptyset, \text{dist}(\tilde{q}, q) \gtrsim R^{1-\delta}\} \lesssim R^{C\delta}. \tag{4-21}$$

Let  $\mathbf{z}_0$  be the center of  $q$ . Then, by the definition of  $\mathcal{W}_1^{\times B}(q; v_1, v'_2)$ , it follows that

$$\bigcup_{\omega_1 \in \mathcal{W}_1^{\times B}(q; v_1, v'_2)} \pi_{\omega_1} \subset \Gamma_1^{v_1, v'_2}(CR^{\frac{1}{2}+\delta}) + \mathbf{z}_0.$$

If  $\omega_2 \in \mathcal{W}_2$ , then it follows from Lemma 3.2 that the intersection

$$\pi_{\omega_2} \cap \left( \bigcup_{\omega_1 \in \mathcal{W}_1^{\times B}(q; v_1, v'_2)} \pi_{\omega_1} \right)$$

is contained in a cube of side length  $O(R^{\frac{1}{2}+\delta})$ . Thus, there are at most  $O(R^{C\delta})$  choices of balls  $\tilde{q} \in \mathcal{Q}(\rho_1, \rho_2)$  such that  $(\tilde{q}, \omega_1)$  is contained in the set in (4-21). On the other hand, since  $\text{dist}(\tilde{q}, q) \gtrsim R^{1-C\delta}$ , we have

$$\#\{w_1 \in \mathcal{W}_1^{\times B}(q; v_1, v'_2) : \pi_{\omega_1} \cap R^\delta \tilde{q} \neq \emptyset, \pi_{\omega_1} \cap R^\delta q \neq \emptyset\} \lesssim R^{C\delta}. \tag{4-22}$$

To see this, by scaling it is enough to check that the map  $S_1 \ni v \mapsto \sum_{i=1}^k t_j \nabla \varphi_i(v)$  is one-to-one whenever  $|t| = 1$ . But this follows from the condition (1-2) if we take  $S_1$  to be small enough. Thus we obtain the claim (4-21). Hence, we also have (4-9), which finishes the proof of Proposition 4.1. This completes the proof of Theorem 1.1.  $\square$

*Proof of Theorem 1.2.* Thanks to Lemma 3.3, the line of argument in the proof of Theorem 1.1 works without modification except that we need to show (4-22). However, to prove (4-22) we don't need to show  $S_1 \ni v \mapsto \sum_{i=1}^k t_j \nabla \varphi_i(v)$  is one-to-one. Instead, as is clear after rescaling, it is enough to show that  $\Pi^{v_1, v'_2} \ni v \mapsto \sum_{i=1}^k t_j \nabla \varphi_i(v)$  is one-to-one. Let  $t_1, \dots, t_{d-k}$  be a set of vectors spanning the tangent space of  $\Pi^{v_1, v'_2}$  at  $v_0$ . Then the above follows if we show that the matrix

$$(t_1^t, \dots, t_{d-k}^t) \left( \sum_{i=1}^k t_j H \varphi_i(v_0) \right)$$

has rank  $d - k$  for  $|t| = 1$ . In fact,  $t_1, \dots, t_{d-k}$  are almost normal to the span of  $\{\nabla \varphi_i(v_2) - \nabla \varphi_i(v_1) : i = 1, \dots, k\}$ . These vectors are close to  $n_1, \dots, n_{d-k}$ . Hence, assuming that  $S_1$  and  $S_2$  are small enough, the above follows if we show  $N(v_2, v_1) \sum_{i=1}^k t_j H \varphi_i(v_0)$  has rank  $d - k$ . This clearly follows from (1-5). □

### 5. Restriction estimates for complex surfaces

In this section we provide the proofs of Corollary 1.5 and Theorem 1.6. In what follows we set  $k = 2$ ,  $d = 2n$ .

*Proof of Corollary 1.5.* Let  $\varphi_1, \varphi_2$  be given by  $\frac{1}{2} z^t D z = \varphi_1 + i \varphi_2$  so that

$$\varphi_1(x, y) = (x^t D x - y^t D y), \quad \varphi_2(x, y) = x^t D y, \quad (x, y) \in \mathbb{R}^n \times \mathbb{R}^n.$$

In order to prove Corollary 1.5 we need only to show that the condition (1-13) implies the assumptions in Theorem 1.1.

Let us set  $z_j = x_j + i y_j \in \mathbb{C}^n$  for  $j = 1, 2$ ,  $\delta_x = x_2 - x_1$ , and  $\delta_y = y_2 - y_1$ . Then a computation shows that the associated matrix  $M(t, z_1, z_2, z)$  is given by

$$M(t, z_1, z_2, z) = \begin{pmatrix} 0 & 0 & \delta_x^t D & -\delta_y^t D \\ 0 & 0 & \delta_y^t D & \delta_x^t D \\ D \delta_x & D \delta_y & t_1 D & t_2 D \\ -D \delta_y & D \delta_x & t_2 D & -t_1 D \end{pmatrix}.$$

Note that

$$\sum_{j=1}^2 t_j H \varphi_j = \begin{pmatrix} t_1 D & t_2 D \\ t_2 D & -t_1 D \end{pmatrix}.$$

Then, it is easy to see that the inverse of  $\sum_{j=1}^2 t_j H \varphi_j$  is

$$(t_1^2 + t_2^2)^{-1} \begin{pmatrix} t_1 D^{-1} & t_2 D^{-1} \\ t_2 D^{-1} & -t_1 D^{-1} \end{pmatrix}.$$

So, the assumption (1-2) holds. Hence, it suffices to show that (1-13) implies (1-3). By the block matrix formula we only need to check

$$\det \left[ \begin{pmatrix} \delta_x^t D & -\delta_y^t D \\ \delta_y^t D & \delta_x^t D \end{pmatrix} \begin{pmatrix} t_1 D^{-1} & t_2 D^{-1} \\ t_2 D^{-1} & -t_1 D^{-1} \end{pmatrix} \begin{pmatrix} D \delta_x & D \delta_y \\ -D \delta_y & D \delta_x \end{pmatrix} \right] \neq 0.$$

By a direct computation it is not difficult to see that the left-hand side equals<sup>4</sup>

$$-(t_1^2 + t_2^2)((\delta_x^t D\delta_x - \delta_y^t D\delta_y)^2 + 4(\delta_x^t D\delta_y)^2).$$

Since  $(z_2 - z_1)^t D(z_2 - z_1) = \delta_x^t D\delta_x - \delta_y^t D\delta_y + 2i\delta_x^t D\delta_y$ , it is now clear that (1-13) implies (1-3).  $\square$

*Proof of Theorem 1.6.* From the bilinear estimate we can get the linear estimate by adapting the arguments in [Tao et al. 1998; Vargas 2005; Lee 2006]. Since  $D$  is nonsingular and symmetric, by making use of linear transforms we may assume that

$$D = \begin{pmatrix} 1 & 0 \\ 0 & \pm 1 \end{pmatrix},$$

and so we have either  $\Phi(z_1, z_2) = z_1^2 + z_2^2 = (z_1 + iz_2)(z_1 - iz_2)$  or  $\Phi(z_1, z_2) = z_1^2 - z_2^2 = (z_1 + z_2)(z_1 - z_2)$ . By a linear change of variables the problem can be further reduced to showing Theorem 1.6 when  $\Phi(z_1, z_2) = z_1 z_2$ .

The following is an immediate consequence of Theorem 1.1 and the translation invariance of the bilinear estimate.

**Lemma 5.1.** *Let  $\Phi(z_1, z_2) = z_1 z_2$  and  $Q_1, Q_2 \subset \mathbb{C}^2$  be closed cubes. Assume that*

$$2^4 \geq |z_1 - w_1| \geq 2^{-1} \quad \text{and} \quad 2^4 \geq |z_2 - w_2| \geq 2^{-1}$$

*whenever  $(z_1, z_2) \in Q_1$  and  $(w_1, w_2) \in Q_2$ . If  $\text{supp}(f) \subset Q_1$  and  $\text{supp}(g) \subset Q_2$ , then for  $q > \frac{10}{3}$  and  $\frac{1}{p} + \frac{5}{3q} < 1$ ,*

$$\|Ef \, Eg\|_{\frac{q}{2}} \leq C_{p,q} \|f\|_p \|g\|_p.$$

In the next lemma the hypothesis of “nonvanishing rotational curvature” is weakened to the usual separation condition. But then, for the conclusion to hold, the pair  $(\frac{1}{p}, \frac{1}{q})$  needs to satisfy a more restrictive condition. This lemma is an analog of Proposition 4.1 in [Lee 2006].

**Lemma 5.2.** *Let  $Q_1, Q_2$  be closed cubes in  $\mathbb{C}^2$  such that  $\text{dist}(Q_1, Q_2) \geq 1$ . If  $\text{supp}(f) \subset Q_1$  and  $\text{supp}(g) \subset Q_2$ , then there is a constant  $C_{p,q}$  such that*

$$\|Ef \, Eg\|_{\frac{q}{2}} \leq C_{p,q} \|f\|_p \|g\|_p \quad \text{if } \frac{1}{p} + \frac{2}{q} < 1, \quad q > \frac{10}{3},$$

or

$$\|E\chi_F \, E\chi_G\|_{\frac{q}{2}} \lesssim \|f\|_{p,1} \|g\|_{p,1} \quad \text{if } \frac{1}{p} + \frac{2}{q} = 1, \quad q > \frac{10}{3}.$$

By translation it is clear that in Lemmas 5.1 and 5.2 the same estimate holds with  $Q_1, Q_2$  replaced by  $Q_1 + a, Q_2 + a$ , respectively, for any  $a \in \mathbb{C}^2$ . It is possible to prove the strong-type estimate  $\|E\chi_F \, E\chi_G\|_{q/2} \lesssim \|f\|_p \|g\|_p$  for  $\frac{1}{p} + \frac{2}{q} = 1, q > \frac{10}{3}$  by making use of the asymmetric estimates which are obtained in the course of proof of Proposition 1.3 and the bilinear interpolation; see, e.g., [Bergh and Löfström 1976, Section 3.13, 5(b)]. However, we have decided not to include the details here, because it does not seem to have any consequences for linear estimates.

<sup>4</sup>In fact, the product of the three matrices is equal to  $\begin{pmatrix} t_1 & -t_2 \\ t_2 & t_1 \end{pmatrix} \begin{pmatrix} \delta_x^t D\delta_x - \delta_y^t D\delta_y & 2\delta_x^t D\delta_y \\ 2\delta_x^t D\delta_y & \delta_y^t D\delta_y - \delta_x^t D\delta_x \end{pmatrix}$ .



*Proof of Lemma 5.2.* By interpolation it suffices to consider the case  $\frac{10}{3} < q \leq 4$  and  $p \leq q$ . By decomposition of the domains, followed by translation and scaling, we may assume that  $Q_1 = H_1 \times K$  and  $Q_2 = H_2 \times K$ , where  $\text{dist}(H_1, H_2) \geq 2^{-1}$  and  $K$  is the unit cube in  $\mathbb{C}$ , centered at the origin.

By a Whitney decomposition, we get

$$(K \times K) \setminus D = \bigcup_{j>1} \bigcup_{\substack{(k,k'): \\ I_k^j \sim I_{k'}^j}} I_k^j \times I_{k'}^j$$

where  $D = \{(z_2, w_2) : z_2 = w_2\}$ , and  $\{I_k^j\}_k$  are the dyadic cubes in  $\mathbb{C}$  of side length  $2^{-j}$ , and as usual the notation  $I_k^j \sim I_{k'}^j$  means that the parent cubes of  $I_k^j$  and  $I_{k'}^j$  are adjacent, while  $I_k^j$  and  $I_{k'}^j$  are not.

Let us set

$$f_k^j(z_1, z_2) = \chi_{I_k^j}(z_2) f(z_1, z_2), \quad g_k^j(w_1, w_2) = \chi_{I_k^j}(w_2) g(w_1, w_2).$$

Then, since the cubes  $I_k^j \times I_{k'}^j$  are almost disjoint, we may write

$$Ef Eg = \sum_j \sum_{\substack{(k,k'): \\ I_k^j \sim I_{k'}^j}} E(f_k^j) E(g_{k'}^j).$$

Since  $q > \frac{10}{3}$ , we get

$$\|Ef Eg\|_{\frac{q}{2}} \leq \sum_j \left\| \sum_{I_k^j \sim I_{k'}^j} E(f_k^j) E(g_{k'}^j) \right\|_{\frac{q}{2}} \lesssim \sum_j \left( \sum_{I_k^j \sim I_{k'}^j} \|E(f_k^j) E(g_{k'}^j)\|_{\frac{q}{2}} \right)^{\frac{2}{q}},$$

where the last inequality follows from Lemma 6.1 in [Tao et al. 1998]. Here, we used the fact that for each fixed  $j$  the supports of the Fourier transforms of  $E(f_k^j) E(g_{k'}^j)$  have uniformly bounded overlap as  $(k, k')$  varies, provided that  $I_k^j \sim I_{k'}^j$ . This is a consequence of the Whitney decomposition. We now claim that if  $I_k^j \sim I_{k'}^j$ , then

$$\|E(f_k^j) E(g_{k'}^j)\|_{\frac{q}{2}} \lesssim 2^{4j(\frac{1}{p} + \frac{2}{q} - 1)} \|f_k^j\|_p \|g_{k'}^j\|_p \tag{5-1}$$

when  $\frac{1}{p} + \frac{5}{3q} < 1$ ,  $q > \frac{10}{3}$ . This is an easy consequence of a translated version of Lemma 5.1. Assuming this for the moment, we will finish the proof. Since  $q \geq p$ , for  $\frac{1}{p} + \frac{5}{3q} < 1$ ,  $4 > q > \frac{10}{3}$ , we have

$$\begin{aligned} \|Ef Eg\|_{\frac{q}{2}} &\leq \sum_j 2^{4j(\frac{1}{p} + \frac{2}{q} - 1)} \left( \sum_{I_k^j \sim I_{k'}^j} \|f_k^j\|_p^{\frac{q}{2}} \|g_{k'}^j\|_p^{\frac{q}{2}} \right)^{\frac{2}{q}} \\ &\lesssim \sum_j 2^{4j(\frac{1}{p} + \frac{2}{q} - 1)} \left( \sum_{I^j} \|f_k^j\|_p^p \right)^{\frac{1}{p}} \left( \sum_{J^j} \|g_{k'}^j\|_p^p \right)^{\frac{1}{p}} \lesssim \sum_j 2^{4j(\frac{1}{p} + \frac{2}{q} - 1)} \|f\|_p \|g\|_p. \end{aligned}$$

Now take  $f = \chi_F$  and  $g = \chi_G$  for measurable sets  $F, G$  contained in  $V_1, V_2$ , respectively. Fix  $p, q$  with  $4 > q > \frac{10}{3}$ ,  $\frac{1}{p} + \frac{2}{q} = 1$ , and choose  $p_1$  and  $p_2$  such that  $\frac{1}{p_j} + \frac{5}{3q} < 1$ ,  $j = 1, 2$ , and

$$\frac{1}{p_1} + \frac{2}{q} - 1 = \eta, \quad \frac{1}{p_2} + \frac{2}{q} - 1 = -\eta$$

for some small  $\eta > 0$ . Then by applying the last estimate for  $p = p_1$  and  $p = p_2$ , we obtain

$$\begin{aligned} \|E\chi_F E\chi_G\|_{\frac{q}{2}} &\lesssim \sum_j \min\{2^{4j\eta}|F|^{\eta+1-\frac{2}{q}}|G|^{\eta+1-\frac{2}{q}}, 2^{-4j\eta}|F|^{-\eta+1-\frac{2}{q}}|G|^{-\eta+1-\frac{2}{q}}\} \\ &\lesssim |F|^{1-\frac{2}{q}}|G|^{1-\frac{2}{q}} = |F|^{1/p}|G|^{\frac{1}{p}}. \end{aligned}$$

This shows the estimate  $\|E\chi_F E\chi_G\|_{q/2} \lesssim \|f\|_{p,1} \|g\|_{p,1}$  for  $\frac{1}{p} + \frac{2}{q} = 1$ ,  $4 > q > \frac{10}{3}$ .

Now it remains to show (5-1). Clearly,  $I_k^j$  and  $I_{k'}^j$  are contained in a ball of radius  $2^{2-j}$  and  $\text{dist}(I_k^j, I_{k'}^j) \geq 2^{-1-j}$ . Hence, by a change of variables,

$$\begin{aligned} E(f_k^j)(w) &= 2^{-2j} E(f_k^j(\cdot, 2^{-j}\cdot))(w_1, 2^{-j}w_2, 2^{-j}w_3), \\ E(g_{k'}^j)(w) &= 2^{-2j} E(g_{k'}^j(\cdot, 2^{-j}\cdot))(w_1, 2^{-j}w_2, 2^{-j}w_3). \end{aligned}$$

Then we see that  $\text{supp } f_k^j(\cdot, 2^{-j}\cdot) \subset H_1 \times \tilde{I}_1$  and  $\text{supp } g_{k'}^j(\cdot, 2^{-j}\cdot) \subset H_2 \times \tilde{I}_2$  if  $\text{dist}(\tilde{I}_1, \tilde{I}_2) \geq 2^{-1}$  and  $\tilde{I}_1, \tilde{I}_2$  are contained in a ball of radius  $\leq 2^3$ . The assumption of Lemma 5.1 is satisfied with  $f = f_k^j(\cdot, 2^{-j}\cdot)$  and  $g = g_{k'}^j(\cdot, 2^{-j}\cdot)$ . Hence we may apply it to  $E(f_k^j(\cdot, 2^{-j}\cdot))E(g_{k'}^j(\cdot, 2^{-j}\cdot))$  and get (5-1).  $\square$

Once Lemma 5.2 is established, the usual argument in [Tao et al. 1998], used to deduce linear estimates from bilinear ones, works without modification. We omit the details.  $\square$

### Acknowledgements

We thank Andreas Seeger for bringing the reference [Alvarez 1997] to our attention. We would also like to thank the anonymous referee for the reference [Banner 2002].

### References

- [Alvarez 1997] D. Alvarez, *Bounds for some Kakeya-type maximal functions*, Ph.D. thesis, University of California, Berkeley, 1997, available at <https://search.proquest.com/docview/304343467>. MR
- [Bak and Ham 2014] J.-G. Bak and S. Ham, “Restriction of the Fourier transform to some complex curves”, *J. Math. Anal. Appl.* **409**:2 (2014), 1107–1127. MR Zbl
- [Bak and Lee 2004] J.-G. Bak and S. Lee, “Restriction of the Fourier transform to a quadratic surface in  $\mathbb{R}^n$ ”, *Math. Z.* **247**:2 (2004), 409–422. MR Zbl
- [Bak and Seeger 2011] J.-G. Bak and A. Seeger, “Extensions of the Stein–Tomas theorem”, *Math. Res. Lett.* **18**:4 (2011), 767–781. MR Zbl
- [Bak et al. 2002] J.-G. Bak, D. M. Oberlin, and A. Seeger, “Two endpoint bounds for generalized Radon transforms in the plane”, *Rev. Mat. Iberoamericana* **18**:1 (2002), 231–247. MR Zbl
- [Bak et al. 2009] J.-G. Bak, D. M. Oberlin, and A. Seeger, “Restriction of Fourier transforms to curves and related oscillatory integrals”, *Amer. J. Math.* **131**:2 (2009), 277–311. MR Zbl
- [Bak et al. 2013] J.-G. Bak, D. M. Oberlin, and A. Seeger, “Restriction of Fourier transforms to curves: an endpoint estimate with affine arclength measure”, *J. Reine Angew. Math.* **682** (2013), 167–205. MR Zbl
- [Banner 2002] A. D. Banner, *Restriction of the Fourier transform to quadratic submanifolds*, Ph.D. thesis, Princeton University, 2002, available at <https://search.proquest.com/docview/305532402>. MR
- [Bennett et al. 2006] J. Bennett, A. Carbery, and T. Tao, “On the multilinear restriction and Kakeya conjectures”, *Acta Math.* **196**:2 (2006), 261–302. MR Zbl

- [Bergh and Löfström 1976] J. Bergh and J. Löfström, *Interpolation spaces: an introduction*, Grundlehren der Mathematischen Wissenschaften **223**, Springer, 1976. MR Zbl
- [Bourgain and Guth 2011] J. Bourgain and L. Guth, “Bounds on oscillatory integral operators based on multilinear estimates”, *Geom. Funct. Anal.* **21**:6 (2011), 1239–1295. MR Zbl
- [Christ 1982] M. Christ, *Restriction of the Fourier transform to submanifolds of low codimension*, Ph.D. thesis, The University of Chicago, 1982, available at <https://search.proquest.com/docview/303090254>. MR
- [Christ 1985] M. Christ, “On the restriction of the Fourier transform to curves: endpoint results and the degenerate case”, *Trans. Amer. Math. Soc.* **287**:1 (1985), 223–238. MR Zbl
- [Greenleaf and Seeger 2002] A. Greenleaf and A. Seeger, “Oscillatory and Fourier integral operators with degenerate canonical relations”, *Publ. Mat.* extra volume (2002), 93–141. MR Zbl
- [Guth 2016] L. Guth, “A restriction estimate using polynomial partitioning”, *J. Amer. Math. Soc.* **29**:2 (2016), 371–413. MR Zbl
- [Lee 2006] S. Lee, “Bilinear restriction estimates for surfaces with curvatures of different signs”, *Trans. Amer. Math. Soc.* **358**:8 (2006), 3511–3533. MR Zbl
- [Mockenhaupt 1996] G. Mockenhaupt, *Bounds in Lebesgue spaces of oscillatory integral operators*, Habilitation thesis, Universität Siegen, 1996, available at <https://tinyurl.com/Mockenhaupt-thesis>. Zbl
- [Moyua et al. 1999] A. Moyua, A. Vargas, and L. Vega, “Restriction theorems and maximal operators related to oscillatory integrals in  $\mathbb{R}^3$ ”, *Duke Math. J.* **96**:3 (1999), 547–574. MR Zbl
- [Oberlin 2005] D. M. Oberlin, “A restriction theorem for a  $k$ -surface in  $\mathbb{R}^n$ ”, *Canad. Math. Bull.* **48**:2 (2005), 260–266. MR Zbl
- [Stein 1993] E. M. Stein, *Harmonic analysis: real-variable methods, orthogonality, and oscillatory integrals*, Princeton Mathematical Series **43**, Princeton University Press, 1993. MR Zbl
- [Stovall 2016] B. Stovall, “Uniform estimates for Fourier restriction to polynomial curves in  $\mathbb{R}^d$ ”, *Amer. J. Math.* **138**:2 (2016), 449–471. MR Zbl
- [Tao 2003] T. Tao, “A sharp bilinear restrictions estimate for paraboloids”, *Geom. Funct. Anal.* **13**:6 (2003), 1359–1384. MR Zbl
- [Tao and Vargas 2000] T. Tao and A. Vargas, “A bilinear approach to cone multipliers, I: Restriction estimates”, *Geom. Funct. Anal.* **10**:1 (2000), 185–215. MR Zbl
- [Tao et al. 1998] T. Tao, A. Vargas, and L. Vega, “A bilinear approach to the restriction and Keakeya conjectures”, *J. Amer. Math. Soc.* **11**:4 (1998), 967–1000. MR Zbl
- [Vargas 2005] A. Vargas, “Restriction theorems for a surface with negative curvature”, *Math. Z.* **249**:1 (2005), 97–111. MR Zbl
- [Wolff 2001] T. Wolff, “A sharp bilinear cone restriction estimate”, *Ann. of Math. (2)* **153**:3 (2001), 661–698. MR Zbl

Received 23 Jan 2017. Revised 2 Jun 2017. Accepted 12 Jul 2017.

JONG-GUK BAK: bak@postech.ac.kr

Department of Mathematics, Pohang University of Science and Technology, Pohang, South Korea

JUNGIN LEE: jungjinlee@unist.ac.kr

Department of Mathematical Sciences, School of Natural Science, Ulsan National Institute of Science and Technology, Ulsan, South Korea

SANGHYUK LEE: shklee@snu.ac.kr

Department of Mathematical Sciences, Seoul National University, Seoul, South Korea



# Analysis & PDE

msp.org/apde

## EDITORS

EDITOR-IN-CHIEF

Patrick Gérard  
patrick.gerard@math.u-psud.fr  
Université Paris Sud XI  
Orsay, France

## BOARD OF EDITORS

Nicolas Burq	Université Paris-Sud 11, France nicolas.burq@math.u-psud.fr	Werner Müller	Universität Bonn, Germany mueller@math.uni-bonn.de
Massimiliano Berti	Scuola Intern. Sup. di Studi Avanzati, Italy berti@sissa.it	Gilles Pisier	Texas A&M University, and Paris 6 pisier@math.tamu.edu
Sun-Yung Alice Chang	Princeton University, USA chang@math.princeton.edu	Tristan Rivière	ETH, Switzerland riviere@math.ethz.ch
Michael Christ	University of California, Berkeley, USA mchrist@math.berkeley.edu	Igor Rodnianski	Princeton University, USA irod@math.princeton.edu
Charles Fefferman	Princeton University, USA cf@math.princeton.edu	Wilhelm Schlag	University of Chicago, USA schlag@math.uchicago.edu
Ursula Hamenstaedt	Universität Bonn, Germany ursula@math.uni-bonn.de	Sylvia Serfaty	New York University, USA serfaty@cims.nyu.edu
Vaughan Jones	U.C. Berkeley & Vanderbilt University vaughan.f.jones@vanderbilt.edu	Yum-Tong Siu	Harvard University, USA siu@math.harvard.edu
Vadim Kaloshin	University of Maryland, USA vadim.kaloshin@gmail.com	Terence Tao	University of California, Los Angeles, USA tao@math.ucla.edu
Herbert Koch	Universität Bonn, Germany koch@math.uni-bonn.de	Michael E. Taylor	Univ. of North Carolina, Chapel Hill, USA met@math.unc.edu
Izabella Laba	University of British Columbia, Canada ilaba@math.ubc.ca	Gunther Uhlmann	University of Washington, USA gunther@math.washington.edu
Gilles Lebeau	Université de Nice Sophia Antipolis, France lebeau@unice.fr	András Vasy	Stanford University, USA andras@math.stanford.edu
Richard B. Melrose	Massachusetts Inst. of Tech., USA rbm@math.mit.edu	Dan Virgil Voiculescu	University of California, Berkeley, USA dvv@math.berkeley.edu
Frank Merle	Université de Cergy-Pontoise, France Frank.Merle@u-cergy.fr	Steven Zelditch	Northwestern University, USA zelditch@math.northwestern.edu
William Minicozzi II	Johns Hopkins University, USA minicozz@math.jhu.edu	Maciej Zworski	University of California, Berkeley, USA zworski@math.berkeley.edu
Clément Mouhot	Cambridge University, UK c.mouhot@dpms.cam.ac.uk		

## PRODUCTION

production@msp.org  
Silvio Levy, Scientific Editor

---

See inside back cover or [msp.org/apde](http://msp.org/apde) for submission instructions.

---

The subscription price for 2017 is US \$265/year for the electronic version, and \$470/year (+\$55, if shipping outside the US) for print and electronic. Subscriptions, requests for back issues from the last three years and changes of subscriber address should be sent to MSP.


---

Analysis & PDE (ISSN 1948-206X electronic, 2157-5045 printed) at Mathematical Sciences Publishers, 798 Evans Hall #3840, c/o University of California, Berkeley, CA 94720-3840, is published continuously online. Periodical rate postage paid at Berkeley, CA 94704, and additional mailing offices.

---

APDE peer review and production are managed by EditFlow® from MSP.

PUBLISHED BY

 **mathematical sciences publishers**  
nonprofit scientific publishing

<http://msp.org/>

© 2017 Mathematical Sciences Publishers

# ANALYSIS & PDE

Volume 10 No. 8 2017

---

Koszul complexes, Birkhoff normal form and the magnetic Dirac operator NIKHIL SVALE	1793
Incompressible immiscible multiphase flows in porous media: a variational approach CLÉMENT CANCÈS, THOMAS O. GALLOUËT and LÉONARD MONSAINGEON	1845
Resonances for symmetric tensors on asymptotically hyperbolic spaces CHARLES HADFIELD	1877
Construction of two-bubble solutions for the energy-critical NLS JACEK JENDREJ	1923
Bilinear restriction estimates for surfaces of codimension bigger than 1 JONG-GUK BAK, JUNGJIN LEE and SANGHYUK LEE	1961
Complete embedded complex curves in the ball of $\mathbb{C}^2$ can have any topology ANTONIO ALARCÓN and JOSIP GLOBEVNIK	1987
Finite-time degeneration of hyperbolicity without blowup for quasilinear wave equations JARED SPECK	2001
Dimension of the minimum set for the real and complex Monge–Ampère equations in critical Sobolev spaces TRISTAN C. COLLINS and CONNOR MOONEY	2031



2157-5045(2017)10:8;1-Q